

TABLE III. The half-width of $J=3, K=3$ line in the inversion spectrum of ammonia.

Pressure 10^{-2} mm Hg	A obs. Mc/sec.	A calc. Mc/sec.
5	1.27	1.15
2.5	0.57	0.58
1.5	0.34	0.34

the asymptote of relative motion of the colliding one, $P(J, K, M)$ the probability that a molecule is in J, K, M state, n the number of molecules in unit volume, and $p_i(t)$, $p_f(t)$ the perturbation energies of initial and final states of the absorption process as a function of time t ; α is the function of J, K, M, V , and ρ where J, K , and M represent the rotational state of colliding molecules.

Using (1) we can calculate intermolecular potential in the usual manner, resulting in Table II. In this table $\pm$ means that the sign must be taken corresponding to that of $\epsilon_{J,K} - \epsilon_{J',K'}$, and e_m is the m th eigenvalue of a matrix whose element is⁷

$$(M_j, M_j' | V | M_i, M_i') = \frac{d^2}{R^3} \frac{KK'}{J(J+1)J'(J'+1)} \\ \times \{ -2M_i M_i' \delta(M_i, M_j) \delta(M_i', M_j') \\ \pm \frac{1}{2} [(J - M_i + 1)(J + M_i)(J' - M_i')(J' + M_i' + 1)]^{\frac{1}{2}} \\ \times \delta(M_i, M_j + 1) \delta(M_i', M_j' - 1) \\ \pm \frac{1}{2} [(J + M_i + 1)(J - M_i)(J' + M_i')(J' - M_i' + 1)]^{\frac{1}{2}} \\ \times \delta(M_i, M_j - 1) \delta(M_i', M_j' + 1) \},$$

R being intermolecular distance. Since an absorption line in the inversion spectrum corresponds to $\psi_{+,J,K} \rightarrow \psi_{-,J,K}$ transition, we can calculate α for it by means of the above table. Making some approximation and again resorting to (2), we can obtain values as in Table III, showing good agreement with the experiment. The half-width of other lines can be easily calculated.

In all these calculations Boltzmann distribution was assumed over rotational states from $J=0$ to $J=10$. On statistical weight of each state one may refer to Herzberg's book.²

¹ These results were published in *The Research in Chemical Physics* (in Japanese) 9, 1 (1947); 11, 11 (1948). Details of them will be published in the near future in *J. Phys. Soc. Japan*.

² G. Herzberg, *Infra-Red and Raman Spectra of Polyatomic Molecules* (D. Van Nostrand Company, Inc., New York, 1945), pp. 26, 221.

³ J. H. Van Vleck, *The Theory of Electric and Magnetic Susceptibilities* (Oxford University Press, New York, 1932), p. 186.

⁴ H. Schoeffers, *Physik. Zeits.* 41, 89, 98 (1940).

⁵ O. Stern, *Zeits. f. Physik* 41, 563 (1900).

⁶ W. Good, *Phys. Rev.* 70, 213 (1946).

⁷ H. Margenau and D. T. Warren, *Phys. Rev.* 51, 748 (1935).

Erratum: Theory of High Frequency Discharges.

IV

[*Phys. Rev.* 73, 326 (1948)]

H. MARGENAU

Sloane Physics Laboratory, Yale University, New Haven, Connecticut

IN this paper the original authorship of the similarity principle was given erroneously. I offer my apologies to Dr. R. Holm, who introduced and discussed it in *Physik. Zeits.* 15, 289 (1914).

Primary and Secondary Meson Event in Photographic Emulsions

ADAIR MORRISON AND ERIC PICKUP

Physics Division, National Research Council, Ottawa, Canada

July 29, 1948

THROUGH the courtesy of the British Overseas Airways Corporation in carrying plates for us on consecutive transatlantic flights, we have had Ilford nuclear research emulsions exposed to cosmic radiation over a period of time. Examination of these emulsions has yielded many interesting cosmic-ray events, some involving mesons.

In a total of 3.26 cc of emulsion of various types of loading so far examined and yielding just over 1000 cosmic-ray stars, we have noted one double meson event of the type first described by Lattes, Muirhead, Occhialini, and Powell¹ (π - μ -process) in which the secondary meson lies wholly in the emulsion. This event was found in a C_2 emulsion, boron-loaded, 60 microns thick after processing. The primary meson enters through the surface and passes downward through the emulsion, ending 18 microns above the glass. The secondary meson also travels upward through the emulsion and is scattered downwards about 50 microns from the end of its range.

The ranges of the primary and secondary meson as measured in horizontal projection are 176 and 627 microns, respectively. A profile showing the vertical path of the mesons through the emulsion is given in Fig. 1. The vertical coordinates have been corrected for the shrinkage of the emulsion in processing by multiplying by a factor of 2.5. After correction for vertical travel, the primary and secondary ranges become 213 and 655 microns, respectively. The accuracy of the range measurement is believed to be ± 0.5 percent. The corrected range of the secondary is somewhat longer than the average range, 614 ± 8 microns given by Lattes, Occhialini, and Powell.² A mosaic of photomicrographs of the tracks is shown in Fig. 2.

Identification of the tracks as due to mesons was made by their scattering, which is greater than that of protons of comparable range, and by the grain counts as compared with that of protons. The total number of grains in the primary or π -meson track is 141, and the grain counts for the last 50, 100, and 150 microns are 46, 77, and 106. For the secondary or μ -meson the corresponding figures are 263 grains and 43, 80, and 112 grains (average grain size in the emulsion = 0.4 micron). The distances are corrected for slant in the emulsion as deduced from the profile given above, and are thus the distances along the path of the mesons when the event took place.




FIG. 1. Vertical profile along the tracks of the primary and secondary mesons.