

Nonstatic vacuum strings: Exterior and interior solutions

J. A. Stein-Schabes

Theoretical Astrophysics Group, Fermi National Accelerator Laboratory, Batavia, Illinois 60510

(Received 22 January 1986)

New nonstatic cylindrically symmetric solutions of Einstein's equations are presented. Some of these solutions represent stringlike objects. An exterior vacuum solution is matched to a nonvacuum interior solution for different forms of the energy-momentum tensor. They generalize the standard static string.

I. INTRODUCTION

Grand unified theories were first suggested as a possible scenario for the unification of all forces except gravity. It was soon realized that for these theories to work, the grand symmetry presented in the early Universe had to be somehow broken in order to account for the electroweak and nuclear forces as different fields in the low-energy universe that we observe today. One of the immediate and inescapable consequences of this breaking of the symmetry is the formation of topological defects in the Higgs fields. These "knots" are simply the consequence of mismatches in the Higgs-field orientation between regions that were not causally connected initially but come into contact later on, trapping regions of the false vacuum. These defects can be classified into three disjoint classes¹ according to the topological properties of the group describing them. They can be magnetic monopoles, cosmic strings, or domain walls.

Production and the cosmological consequences of these objects have been thoroughly investigated by many authors.² It has been shown³ that the existence of walls is incompatible with present-day observations of the microwave background radiation. Monopole production has been calculated in several simple models⁴ giving an unacceptable number density. However, when more complicated cosmological models are used then overproduction of monopoles can be reduced by a considerable amount.⁵ Furthermore, if inflation takes place in the early Universe then the monopole problem disappears completely.⁶ Vacuum strings, on the other hand, are much more difficult to rule out. In fact it has been shown¹ that it is possible to accommodate a network of stable strings and still explain observations of the microwave background radiation. Furthermore, cosmic strings might even explain galaxy formation and the large-scale structure of the Universe.

There is, at least in principle, one observational consequence of their existence. Strings act as gravitational lenses, producing focusing and even multiple images of luminous sources in the Universe. This effect was first noted by Vilenkin⁷ and Gott and Alpert,⁸ and later used by Gott⁹ to explain some observations made by Paczynski and Gorski¹⁰ of what is believed to be a triple quasi-stellar object, as just being the split image of a single object.

All these discussions have been based on the model of a

string as a vacuum, static exterior and nonvacuum interior solution of Einstein's equations with cylindrical symmetry. The interior solution describes a fluid of constant density ρ and constant pressure P_z along the direction of the string, with the property that $\rho = -P_z$. The two solutions are then matched at the boundary of the string. The purpose of this paper is to present a new string-type solution and some by-products of this. In our case the string energy density is a dynamical variable $\rho = \rho(r, t)$ with r the cylindrical radius. In Sec. II we will give the metric, the field equations, their solutions, and the matching conditions. Finally, some brief conclusions will be drawn.

II. THE FIELD EQUATIONS

We will start with a metric describing a cylindrically symmetric space-time, with two Killing vectors ∂_z and ∂_ϕ :

$$ds^2 = A^2(r)[dt^2 - B^2(t)dz^2] - D^2(t)[dr^2 + G^2(r)d\phi^2], \quad (1)$$

where r , ϕ , and z are cylindrical coordinates defined in the range $0 \leq r < \infty$; $0 \leq \phi < 2\pi$; $-\infty < z < \infty$; $t \geq 0$. The Einstein field equations will be given (in units where $8\pi G = c = \hbar = 1$) by

$$R_\mu^\nu = T_\mu^\nu - \frac{1}{2}\delta_\mu^\nu T. \quad (2)$$

The energy-momentum tensor for the exterior solution will be $T_{\mu(\text{ext})}^\nu = 0$, while for the interior solution it will have the form prescribed in Ref. 7 (also see Refs. 11 and 12):

$$T_{\mu(\text{int})}^\nu = \mu(t)f(r) \text{diag}(1, 0, 0, 1). \quad (3)$$

The energy-momentum tensor reduces to the standard one if we take $\mu = \mu_0$. Now we can write the field equations as

$$\frac{\ddot{B}}{A^2 B} - \frac{A'^2}{A^2 D^2} + 2\frac{\ddot{D}}{A^2 D} - \frac{A''}{AD^2} - \frac{A'G'}{AGD^2} = 0, \quad (4)$$

$$\frac{A'\dot{D}}{A^3 D} = 0, \quad (5)$$

$$\frac{\ddot{D}}{A^2 D} + \frac{\dot{D}^2}{D^2 A^2} + \frac{\dot{D}\dot{B}}{DBA^2} - 2\frac{A''}{AD^2} - \frac{G''}{GD^2} = T_0^0, \quad (6)$$

$$\frac{\ddot{D}}{A^2 D} + \frac{\dot{D}^2}{D^2 A^2} + \frac{\dot{D}\dot{B}}{DBA^2} - 2\frac{A'G'}{AGD^2} - \frac{G''}{GD^2} = T_0^0, \quad (7)$$

$$\frac{\ddot{B}}{A^2 B} + 2\frac{\dot{B}\dot{D}}{BDA^2} - \frac{A'^2}{A^2 D^2} - \frac{A''}{AD^2} - \frac{A'G'}{AGD^2} = 0, \quad (8)$$

Eq. (4) can immediately be integrated. It contains three different cases (an overdot denotes $\partial/\partial t$ and a prime denotes $\partial/\partial r$):

- (i) $A' = 0 \Rightarrow A = \text{const}$,
- (ii) $\dot{D} = 0 \Rightarrow D = \text{const}$,
- (iii) $A' = 0 = \dot{D} \Rightarrow A, D = \text{const}$.

We will study each case separately.

(i) In this case we will take, without loss of generality, $A = 1$. Then the field equations become

$$\frac{\ddot{B}}{B} + 2\frac{\dot{D}}{D} = 0, \quad (9)$$

$$\frac{\ddot{D}}{D} + \frac{\dot{D}^2}{D^2} + \frac{\dot{D}\dot{B}}{DB} - \frac{G''}{GD^2} = T_0^0, \quad (10)$$

$$\frac{\ddot{B}}{B} + 2\frac{\dot{B}\dot{D}}{BD} = 0. \quad (11)$$

Equations (9) and (11) can be integrated simultaneously for the exterior and interior cases as they are independent of the energy-momentum tensor. The solution is

$$B = B_0 \dot{D}, \quad (12)$$

$$\dot{D}^2 = \kappa_1 - \frac{2}{D}. \quad (13)$$

It is clear from (13) that for $\kappa_1 > 0$ the solution is given by

$$D = \frac{1}{\kappa_1} (1 + \cosh \psi), \quad (14a)$$

$$t = t_0 + \frac{1}{\kappa_1^{3/2}} (\psi + \sinh \psi). \quad (14b)$$

For the exterior solution we can take $T_0^0 = 0$ in which case we get, from (10),

$$D\ddot{D} + \dot{D}^2 + \frac{\dot{B}}{B} D\dot{D} = \kappa_2 = \frac{G''}{G}, \quad (15)$$

where κ_2 is an arbitrary constant. The left-hand side of this triple equality becomes a constraint on the solutions for D and B , which can be easily satisfied provided we choose $\kappa_1 = \kappa_2$, while the right-hand side just yields

$$G'' - \kappa_1 G = 0. \quad (16)$$

Since $\kappa_1 > 0$ the solution of this equation is given by

$$G(r) = C_1 e^{(\kappa_1 r)^{1/2}} + C_2 e^{-(\kappa_1 r)^{1/2}}, \quad (17)$$

with C_1 and C_2 integration constants. In order to find an interior solution we have to specify the form of T_0^0 . The continuity equation $T_{;\nu}^{\mu\nu} = 0$ together with Eq. (10) suggest

$$T_0^0 = \frac{f(r)}{D^2}.$$

Then Eq. (10) becomes a separable equation in its radial and time-dependent parts, so obtaining

$$D\ddot{D} + \dot{D}^2 + \frac{\dot{B}}{B} D\dot{D} = \kappa_3 = \frac{G''}{G} + f(r), \quad (18)$$

where $f(r)$ is an arbitrary function of r . The first equality is satisfied by the same B and D found previously provided we take $\kappa_3 = \kappa_1$. However, the right-hand side now gives a different equation: namely,

$$G'' + [f(r) - \kappa_1]G = 0. \quad (19)$$

This equation is identical to a Schrödinger equation describing a free particle moving in a potential $f(r)$ with energy κ_1 . Clearly, to solve this equation we need to specify the form of $f(r)$. For the solution of (19) to be valid we require $G(r)$ and $G'(r)$ to be continuous at the boundary of the string, i.e.,

$$G_{\text{ext}}(r_*) = G_{\text{int}}(r_*), \quad (20)$$

$$G'_{\text{ext}}(r_*) = G'_{\text{int}}(r_*). \quad (21)$$

This implies that any solution of the one-dimensional Schrödinger equation will be a valid solution of (19) (we are not establishing any physical relation between these two systems but merely pointing out the mathematical similarity). Before attempting to solve (19) for a specific potential, let us examine the form of the metric and the matching conditions.

With the help of Eq. (12) the exterior and interior metrics can be put in the form

$$ds^2 = dt^2 - D^2(t)[dr^2 + G^2(r)d\phi^2] - \dot{D}^2(t)dz^2, \quad (22)$$

where $D(t)$ is given by Eq. (14) and $G(r)$ by Eq. (17) for the exterior and by Eq. (19) for the interior. Since the coordinate systems used to describe the exterior and the interior are the same, then the matching conditions are given by Eqs. (20) and (21). $G(r)$ and $G'(r)$ must be continuous at the boundary. We only need to choose a sensible "potential" for which Eq. (19) has a solution consistent with the matching conditions.

We will now analyze in more detail a specific potential. If we take $f(r) = f_0$ for the interior solution, then (19) has the solution

$$G(r) = \tilde{C}_1 e^{\sqrt{\alpha}r} + \tilde{C}_2 e^{-\sqrt{\alpha}r}, \quad \alpha > 0, \quad (23a)$$

$$G(r) = \tilde{C}_1 \cos(\sqrt{\alpha}r) + \tilde{C}_2 \sin(\sqrt{\alpha}r), \quad \alpha < 0, \quad (23b)$$

with $\alpha \equiv f_0 - \kappa_1$. We shall now concentrate on the second case and will take $\tilde{C}_1 = 0$. Then the metric can be written as

$$ds^2 = dt^2 - D^2(t)[dr^2 + \tilde{C}_2 \sin^2(\sqrt{\alpha}r)d\phi^2] - \dot{D}^2(t)dz^2. \quad (24)$$

For every $t = \text{const}$ hypersurface it is possible to rescale the coordinates so as to get

$$ds^2 = - \left[d\tilde{r}^2 + C_0^2 \sin^2 \left[\frac{\tilde{r}}{r_0} \right] d\phi^2 + d\tilde{z}^2 \right]. \quad (25)$$

This slice is identical to the equivalent slice of the standard string.⁷ The main difference is that for each hypersurface we need to rescale differently in order to obtain the standard string. In this solution the interior is a nonstatic space-time.

For the exterior solution let $C_1 = -C_2 \equiv \frac{1}{2}C$, then

$$G(r) = C \sinh(\sqrt{\kappa_1} r) . \quad (26)$$

Near the string wall (r small) the metric can be approximated by

$$ds^2 = dt^2 - D^2(t)(dr^2 + C^2 r^2 d\phi^2) - \dot{D}^2(t) dz^2 \quad (27)$$

and again a $t = \text{const}$ slice reduces to the equivalent slice of the static exterior solution.

The analysis of the stability of this solution, and those obtained for more complicated "potentials" is underway and will be reported elsewhere. However, we might speculate that these types of solution will be unstable for any potential other than $f(r) = \text{const}$ or $f(r) = \delta(r)$, because if the density distribution is nonhomogeneous then different parts of the string (cross section) would feel different tensions and this would tend to disrupt the string. There might be some gravitational radiation coming from this configuration as it settles into a lower-energy configuration (maybe the static case). There could even be a "Birkhoff-like" theorem for cylindrically symmetric solutions.

(ii) In this case we shall take $D=1$. Now the field equations read

$$\frac{A''}{A} + \frac{A'^2}{A^2} + \frac{A'G'}{AG} - \frac{\ddot{B}}{A^2 B} = 0 , \quad (28)$$

$$2 \frac{A''}{A} + \frac{G''}{G} = -T_0^0 , \quad (29)$$

$$2 \frac{A'G'}{AG} + \frac{G''}{G} = -T_0^0 . \quad (30)$$

If we now substitute (29) into (30) we get

$$\frac{A''}{A} - \frac{A'G'}{AG} = 0 \quad (31)$$

which can be easily integrated to give

$$G = G_0 A' . \quad (32)$$

For the exterior solution we take as before $T_0^0 = 0$ and using (29) and (32) we arrive at

$$\frac{A'''}{A} + 2 \frac{A''}{A} = 0 \quad (33)$$

which immediately integrates to give

$$A(r) = \frac{1}{4}(2C_2 - C_1) + \frac{C_1}{2}r + C_3 e^{-2r} , \quad (34)$$

$$G(r) = G_0 \left[\frac{C_1}{2} - 2C_3 e^{-2r} \right] . \quad (35)$$

Equation (30) is identically satisfied, while (28) now yields the triple equality

$$\left[\frac{A''}{A} + \frac{A'G'}{AG} + \frac{A'^2}{A^2} \right] A^2 = \kappa_1 = \frac{\ddot{B}}{B} . \quad (36)$$

The right-hand side of (36) again becomes a constraint on the form of $A(r)$, which can be satisfied provided we take $C_3 = 0$ and $C_1 = 2\kappa_1^{1/2}$. We can then write $A(r)$ as

$$A(r) = A_0 + \kappa_1^{1/2} r , \quad (37)$$

with $A_0 = \frac{1}{2}(C_2 - \kappa_1^{1/2})$. The left-hand side of (36) yields the following equation for $B(t)$:

$$\ddot{B} - \kappa_1 B = 0 \quad (38)$$

which has the same (formal) solution as (17) if we exchange $G \rightarrow B$ and $r \rightarrow t$.

For the interior case, the solution described by Eq. (32) is still valid, and when substituted into (28), for the case $f(r) = f_0$ we get

$$\frac{A'''}{A} + 2 \frac{A''}{A} = -T_0^0 = -f_0 \equiv -\frac{2g_0}{27} , \quad (39)$$

with g_0 a new constant defined in terms of f_0 . The general solution to Eq. (27) is

$$A(r) = C_1 e^{m_1 r} + e^{ar} [C_2 \cos(br) + C_3 \sin(br)] , \quad (40)$$

with

$$b = \frac{1}{2\sqrt{3}} \{ [-(g_0 + 1) + g_0^{1/2}(g_0 + 2)^{1/2}]^{1/3} + [(g_0 + 1) + g_0^{1/2}(g_0 + 2)^{1/2}]^{1/3} \} ,$$

$$m_1 = \frac{1}{3} \{ [(g_0 + 1) - g_0^{1/2}(g_0 + 2)^{1/2}]^{1/3} + [(g_0 + 1) + g_0^{1/2}(g_0 + 2)^{1/2}]^{1/3} + 1 \} .$$

For small r the solution for $A(r)$ can be approximated by

$$A(r) \simeq C_1 + bC_3 r \quad (41)$$

for which the matching conditions are satisfied provided we take $bC_3 = \kappa_1^{1/2}$ and $C_1 = A_0$. Now we can rewrite the metric in the form

$$ds^2 = A^2(r)[dt^2 - B^2(t)dz^2] - dr^2 - G_0^2 A'^2(r)d\phi^2 . \quad (42)$$

This case does not reduce to the static string case, and even though it has the right Killing vectors and both exterior and interior solutions can be found and matched, it is extremely difficult to give any interpretation to this space-time.

(iii) In this case we take $A=1=D$. For this case the field equations become

$$\frac{\ddot{B}}{B} = 0 , \quad (43)$$

$$\frac{G''}{G} = -T_0^0 . \quad (44)$$

For the exterior solution we get

$$B = B_0 t + B_1 , \quad (45)$$

$$G = G_0 r + G_1 , \quad (46)$$

where B_0 , B_1 , G_0 , and G_1 are integration constants. We shall consider only one, namely, $G_1=0$. Then the metric becomes

$$ds^2 = dt^2 - (B_0 t + B_1)^2 dz^2 - (dr^2 + G_0^2 r^2 d\phi^2), \quad (47)$$

$B(t)$ for the interior is given by Eq. (45), while $G(r)$ is the solution to

$$G'' + f(r)G = 0. \quad (48)$$

For the special case we are interested in, namely, $f(r)=f_0$, we get

$$G(r) = G_1 e^{(f_0 r)^{1/2}} + G_2 e^{-(f_0 r)^{1/2}}. \quad (49)$$

If we take $G_1 = G_2 \equiv \frac{1}{2} G_1$ then for small r we get

$$ds^2 = dt^2 - (B_0 t + B_1)^2 dz^2 - (dr^2 + G_1^2 f_0^2 r^2 d\phi^2). \quad (50)$$

The matching of these solutions is achieved provided we take $G_1 f_0 = G_0$. Two particular cases are worth mentioning. If $B_0=0$ then the solution becomes the standard static string described in the literature. When $B_1=0$ we get a slightly different metric which describes a string being stretched by the horizon as t increases, and whose cross section remains the same for every t hypersurface.

III. CONCLUSIONS

In this paper we have presented some new cylindrically symmetric solutions of Einstein's field equations. Some of these solutions represent stringlike objects for which an interior nonvacuum solution was matched to an exterior vacuum solution. We found that the solutions naturally fall into three classes depending on whether $A(r)$ or $D(t)$ or both are constants. The static string solution described by Vilenkin⁷ and Gott⁹ is a particular case of our solution. The nonstatic behavior of the solution could change some of the results obtained using static strings. The lensing property, for one, will probably be modified as the amplification or reduction of the amplitude due to focusing or defocusing of bundles of light rays coming from the source are changed.

As mentioned earlier, the analysis of the stability of these solutions is still underway. However, the solution where the density distribution is a delta function or a constant ought to be stable.

ACKNOWLEDGMENTS

This work was partially supported by the Department of Energy and NASA. Part of this work was done at the Astronomy Centre, University of Sussex, England.

¹T. W. B. Kibble, J. Phys. A 9, 1382 (1976).

²J. Preskill, in *The Very Early Universe*, edited by G. W. Gibbons, S. W. Hawking, and S. T. C. Siklos (Cambridge University Press, Cambridge, 1983).

³Ya. B. Zeldovich, I. Yu. Kobzarev, and L. B. Okun, Zh. Eksp. Teor. Fiz. 67, 3 (1974) [Sov. Phys. JETP 40, 1 (1975)].

⁴Ya. B. Zeldovich and M. Yu. Khlopov, Phys. Lett. 79B, 239 (1978); J. Preskill, Phys. Rev. Lett. 43, 1365 (1979); for a review, see *Monopole '83*, edited by J. L. Stone (Plenum, New York, 1984).

⁵P. Fargion, Phys. Lett. 127B, 35 (1983); J. Stein-Schabes, Mon. Not. R. Astron. Soc., 216, 809 (1985).

⁶M. S. Turner, Report No. Fermilab-Conf-85/153-A, 1985 (unpublished).

⁷A. Vilenkin, Phys. Rev. D 23, 852 (1981).

⁸J. Gott and M. Alpert, Gen. Relativ. Gravit. 16, 243 (1984).

⁹J. Gott, Astrophys. J. 288, 422 (1985).

¹⁰B. Paczynski and K. Gorski, Astrophys. J. 248, L101 (1981).

¹¹A. Vilenkin, Phys. Rep. 121, 263 (1985).

¹²D. Garfinkle, Phys. Rev. D 32, 1323 (1985).