

Search for an exotic $S = -2$, $Q = -2$ baryon resonance at a mass near 1862 MeV in quasisreal photoproduction

A. Airapetian,¹⁸ N. Akopov,³⁰ Z. Akopov,³⁰ M. Amarian,^{8,30} A. Andrus,¹⁶ E. C. Aschenauer,⁸ W. Augustyniak,²⁹ R. Avakian,³⁰ A. Avetissian,³⁰ E. Avetissian,¹² P. Bailey,¹⁶ D. Balin,²¹ M. Beckmann,⁷ S. Belostotski,²¹ N. Bianchi,¹² H. P. Blok,^{20,28} H. Böttcher,⁸ A. Borissov,¹⁵ A. Borysenko,¹² M. Bouwhuis,¹⁶ A. Brüll,¹⁷ V. Bryzgalov,²² M. Capiluppi,¹¹ G. P. Capitani,¹² T. Chen,⁴ G. Ciullo,¹¹ M. Contalbrigo,¹¹ P. F. Dalpiaz,¹¹ W. Deconinck,¹⁸ R. De Leo,³ M. Demey,²⁰ L. De Nardo,¹ E. De Sanctis,¹² E. Devitsin,¹⁹ M. Diefenthaler,¹⁰ P. Di Nezza,¹² J. Drechsler,²⁰ M. Düren,¹⁴ M. Ehrenfried,¹⁰ A. Elalaoui-Moulay,² G. Elbakian,³⁰ F. Ellinghaus,⁶ U. Elschenbroich,¹³ R. Fabbri,²⁰ A. Fantoni,¹² L. Felawka,²⁶ S. Frullani,²⁴ A. Funel,¹² G. Gapienko,²² V. Gapienko,²² F. Garibaldi,²⁴ K. Garrow,²⁶ G. Gavrilov,^{7,26} V. Gharibyan,³⁰ F. Giordano,¹¹ O. Grebeniuk,²¹ I. M. Gregor,⁸ C. Hadjidakis,¹² K. Hafidi,² M. Hartig,¹⁴ D. Hasch,¹² W. H. A. Hesselink,^{20,28} A. Hillenbrand,¹⁰ M. Hoek,¹⁴ Y. Holler,⁷ B. Hommez,¹³ I. Hristova,⁸ G. Iarygin,⁹ A. Ivanilov,²² A. Izotov,²¹ H. E. Jackson,² A. Jgoun,²¹ R. Kaiser,¹⁵ T. Keri,¹⁴ E. Kinney,⁶ A. Kisselev,^{6,21} T. Kobayashi,²⁷ M. Kopytin,⁸ V. Korotkov,²² V. Kozlov,¹⁹ B. Krauss,¹⁰ V. G. Krivokhijine,⁹ L. Lagamba,³ L. Lapikás,²⁰ A. Laziev,^{20,28} P. Lenisa,¹¹ P. Liebing,⁸ L. A. Linden-Levy,¹⁶ W. Lorenzon,¹⁸ H. Lu,⁵ J. Lu,²⁶ S. Lu,¹⁴ B.-Q. Ma,⁴ B. Maiheu,¹³ N. C. R. Makins,¹⁶ Y. Mao,⁴ B. Marianski,²⁹ H. Marukyan,³⁰ F. Masoli,¹¹ V. Mexner,²⁰ N. Meyners,⁷ T. Michler,¹⁰ O. Mikloukho,²¹ C. A. Miller,^{1,26} Y. Miyachi,²⁷ V. Muccifora,¹² M. Murray,¹⁵ A. Nagaitsev,⁹ E. Nappi,³ Y. Naryshkin,²¹ M. Negodaev,⁸ W.-D. Nowak,⁸ K. Oganessyan,^{7,12} H. Ohsuga,²⁷ A. Osborne,¹⁵ N. Pickert,¹⁰ D. H. Potterveld,² M. Raithel,¹⁰ D. Reggiani,¹⁰ P. E. Reimer,² A. Reischl,²⁰ A. R. Reolon,¹² C. Riedl,¹⁰ K. Rith,¹⁰ G. Rosner,¹⁵ A. Rostomyan,³⁰ L. Rubacek,¹⁴ J. Rubin,¹⁶ D. Ryckbosch,¹³ Y. Salomatin,²² I. Sanjiev,^{2,21} I. Savin,⁹ A. Schäfer,²³ G. Schnell,²⁷ K. P. Schüller,⁷ J. Seele,⁶ R. Seidl,¹⁰ B. Seitz,¹⁴ C. Shearer,¹⁵ T.-A. Shibata,²⁷ V. Shutov,⁹ K. Sinram,⁷ W. Sommer,¹⁴ M. Stancari,¹¹ M. Statera,¹¹ E. Steffens,¹⁰ J. J. M. Steijger,²⁰ H. Stenzel,¹⁴ J. Stewart,⁸ F. Stinzinger,¹⁰ P. Tait,¹⁰ H. Tanaka,²⁷ S. Taroian,³⁰ B. Tchuiko,²² A. Terkulov,¹⁹ A. Trzcinski,²⁹ M. Tytgat,¹³ A. Vandenbroucke,¹³ P. B. van der Nat,²⁰ G. van der Steenhoven,²⁰ Y. van Haarlem,¹³ V. Vikhrov,²¹ M. G. Vincet,¹ C. Vogel,¹⁰ J. Volmer,⁸ S. Wang,⁵ J. Wendland,^{25,26} Y. Ye,⁵ Z. Ye,⁷ S. Yen,²⁶ B. Zihlmann,¹³ and P. Zupranski²⁹

(HERMES Collaboration)

¹Department of Physics, University of Alberta, Edmonton, Alberta, Canada T6G 2J1

²Physics Division, Argonne National Laboratory, Argonne, Illinois 60439-4843, USA

³Istituto Nazionale di Fisica Nucleare, Sezione di Bari, 70124 Bari, Italy

⁴School of Physics, Peking University, Beijing 100871, China

⁵Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China

⁶Nuclear Physics Laboratory, University of Colorado, Boulder, Colorado 80309-0446, USA

⁷DESY, 22603 Hamburg, Germany

⁸DESY, 15738 Zeuthen, Germany

⁹Joint Institute for Nuclear Research, 141980 Dubna, Russia

¹⁰Physikalisches Institut, Universität Erlangen-Nürnberg, 91058 Erlangen, Germany

¹¹Istituto Nazionale di Fisica Nucleare, Sezione di Ferrara and Dipartimento di Fisica, Università di Ferrara, 44100 Ferrara, Italy

¹²Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, 00044 Frascati, Italy

¹³Department of Subatomic and Radiation Physics, University of Gent, 9000 Gent, Belgium

¹⁴Physikalisches Institut, Universität Gießen, 35392 Gießen, Germany

¹⁵Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, United Kingdom

¹⁶Department of Physics, University of Illinois, Urbana, Illinois 61801-3080, USA

¹⁷Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

¹⁸Randall Laboratory of Physics, University of Michigan, Ann Arbor, Michigan 48109-1120, USA

¹⁹Lebedev Physical Institute, 117924 Moscow, Russia

²⁰Nationaal Instituut voor Kernfysica en Hoge-Energiefysica (NIKHEF), 1009 DB Amsterdam, The Netherlands

²¹Petersburg Nuclear Physics Institute, St. Petersburg, Gatchina, 188350 Russia

²²Institute for High Energy Physics, Protvino, Moscow region, 142281 Russia

²³Institut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany

²⁴Istituto Nazionale di Fisica Nucleare, Sezione Roma 1, Gruppo Sanità and Physics Laboratory,

Istituto Superiore di Sanità, 00161 Roma, Italy

²⁵Department of Physics, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6

²⁶TRIUMF, Vancouver, British Columbia, Canada V6T 2A3

²⁷*Department of Physics, Tokyo Institute of Technology, Tokyo 152, Japan*²⁸*Department of Physics and Astronomy, Vrije Universiteit, 1081 HV Amsterdam, The Netherlands*²⁹*Andrzej Soltan Institute for Nuclear Studies, 00-689 Warsaw, Poland*³⁰*Yerevan Physics Institute, 375036 Yerevan, Armenia*

(Received 10 December 2004; published 10 February 2005)

A search for an exotic baryon resonance with $S = -2$, $Q = -2$ has been performed in quasireal photoproduction on a deuterium target through the decay channel $\Xi^- \pi^- \rightarrow \Lambda \pi^- \pi^- \rightarrow p \pi^- \pi^- \pi^-$. No evidence for a previously reported $\Xi^{--}(1860)$ resonance is found in the $\Xi^- \pi^-$ invariant mass spectrum. An upper limit for the photoproduction cross section of 2.1 nb is found at the 90% confidence level. The photoproduction cross section for the $\Xi^0(1530)$ is found to be between 9 and 24 nb.

DOI: 10.1103/PhysRevD.71.032004

PACS numbers: 12.39.Mk, 13.60.-r, 13.60.Rj, 14.20.-c

The prediction for the existence of narrow exotic baryon resonances [1], based on the chiral soliton model, has triggered an intensive search for the exotic members of an antidecuplet with spin 1/2. In this antidecuplet [1–3] all three vertices are manifestly exotic. The lightest exotic member of this antidecuplet lying at its apex, named the Θ^+ , was predicted [1] to have a mass of 1530 MeV and a narrow width. It corresponds to a $uudd\bar{s}$ configuration and decays through the channels $\Theta^+ \rightarrow pK^0$ or $\Theta^+ \rightarrow nK^+$.

Experimental evidence for the Θ^+ first came from the observation of a narrow resonance at $1540 \pm 10(\text{syst})$ MeV in the K^- missing mass spectrum for the $\gamma n \rightarrow K^+ K^- n$ reaction on ^{12}C [4]. The decay mode corresponds to a $S = +1$ resonance and signals an exotic pentaquark state with quark content ($uudd\bar{s}$). Further evidence emerged from a series of experiments, with the observation of narrow peaks [5–15] in the nK^+ and pK_S^0 invariant mass spectra near 1530 MeV, in most cases with a width that is limited by the experimental resolution. Some doubts have been raised recently, however, because of potential experimental artifacts [16] and the failure to observe a signal in other experiments [17–23].

Experimental evidence for a second exotic member of the antidecuplet came from the reported observation of a $S = -2$, $Q = -2$ baryon resonance in proton-proton collisions at $\sqrt{s} = 17.2$ GeV at the CERN SPS [24]. A narrow peak at a mass of about 1862 MeV in the $\Xi^- \pi^-$ invariant mass spectrum is proposed as a candidate for the predicted exotic $\Xi_{3/2}^{--}$ baryon with $S = -2$, $I = \frac{3}{2}$ and a quark content of ($ddss\bar{u}$). At the same mass, a peak is observed that is a candidate for the $\Xi_{3/2}^0$ member of this isospin quartet. The corresponding antibaryon spectra show enhancements at the same invariant mass. The observed mass of 1862 MeV falls below the prediction of Ref. [1] and above a prediction of Ref. [25], although closer to the latter. However, the result of Ref. [24] has been disputed [26]. In addition, this resonance has not been confirmed by other experimental searches [22,27,28]. Many further searches for the Ξ^{--} are presently underway [29] for which no final results are available yet.

This paper presents the results of a search for the Ξ^{--} in quasireal photoproduction on deuterium. The data were

obtained by the HERMES experiment with the 27.6 GeV positron beam of the HERA storage ring at DESY. Stored beam currents ranged from 9 to 45 mA. An integrated luminosity of 296 pb^{-1} was collected on a deuterium gas target. The target was either longitudinally polarized (223 pb^{-1}) or unpolarized (73 pb^{-1}). The yields were summed over two spin orientations when the target was polarized.

The HERMES spectrometer [30] consists of two identical halves located above and below the positron beam pipe and has an angular acceptance of ± 170 mrad horizontally, and $\pm(40\text{--}140)$ mrad vertically. The trigger was formed by either a coincidence between scintillator hodoscopes, a preshower detector and a lead-glass calorimeter, or a coincidence between three scintillator hodoscopes and two tracking planes, requiring that at least one charged track appears in each of the detector halves of the spectrometer.

The analysis searched for inclusive photoproduction of a Ξ^{--} followed by the decay $\Xi^{--} \rightarrow \Xi^- \pi^- \rightarrow \Lambda \pi^- \pi^- \rightarrow p \pi^- \pi^- \pi^-$ or a $\Xi^0 \rightarrow \Xi^- \pi^+ \rightarrow \Lambda \pi^- \pi^+ \rightarrow p \pi^- \pi^- \pi^+$. Events selected contained at least four tracks: three charged pions in coincidence with one proton. Identification of charged pions and protons was accomplished with a ring-imaging Čerenkov (RICH) detector [31] which provides separation of pions, kaons, and protons over most of the kinematic acceptance of the spectrometer. The protons were restricted to a momentum range of 2–15 GeV/c and pions to a range of 0.25–15 GeV/c.

Based on the intrinsic tracking resolution, the required event topology included a minimum distance of approach between a proton and a negative pion track less than 1.5 cm (the midpoint of which is defined as the Λ decay vertex), a minimum distance of approach between a second negative pion and the reconstructed Λ track less than 1.0 cm (the midpoint of which is defined as the Ξ^- decay vertex), a minimum distance of approach between a third pion and the reconstructed Ξ^- track less than 2.5 cm (the midpoint of which is defined as the production vertex), a radial distance of the production vertex from the positron beam axis less than 6 mm, a z coordinate of the production vertex in the ± 20 cm long target cell within $-18 < z < +18$ cm along the beam direction, a Λ decay length (separation of Ξ^- and Λ decay vertices) greater than 7 cm, and a Ξ^-

decay length (separation of production and Ξ^- decay vertices) greater than 10 cm. Both possible $\pi^+\pi^-$ combinations were examined for a K_S^0 peak in the $M_{\pi^+\pi^-}$ mass spectrum, and none were found.

The first step in the analysis was to search for Λ candidates. The resulting invariant $M_{p\pi^-}$ spectrum is shown in Fig. 1(a). The position of the observed Λ peak at $1115.73 \pm 0.01(\text{stat})$ MeV is in good agreement with the nominal value [32]. Events were selected with a $M_{p\pi^-}$ invariant mass within $\pm 3\sigma$ of the centroid of the Λ peak. They were then combined with a π^- to form the Ξ^- candidates. The resulting invariant $M_{\Lambda\pi^-}$ spectrum, shown in Fig. 1(b), yields a Ξ^- peak at $1321.8 \pm 0.3(\text{stat})$ MeV, which is in good agreement with the expected value of 1321.3 MeV [32].

The next step was to search for $\Xi_{3/2}^{--}$ ($\Xi_{3/2}^0$) candidates in the $M_{\Xi^-\pi^-}$ ($M_{\Xi^-\pi^+}$) spectra. Events were selected with a $M_{\Lambda\pi^-}$ invariant mass within $\pm 3\sigma$ of the centroid of the Ξ^- peak. The resulting spectrum of the invariant mass of the $p\pi^-\pi^-\pi^-$ system is displayed in Fig. 2. No peak structure is observed near 1862 MeV. Figure 3 shows the resulting spectrum of the invariant mass of the $p\pi^-\pi^-\pi^+$ system. While no peak structure is observed near 1862 MeV, one appears at the mass of the known $\Xi^0(1530)$.

The shape of the background in Fig. 2 (Fig. 3) was determined from randomly mixing Ξ^- events with π^- (π^+) tracks from other events. Each resulting distribution was fit with a broad Gaussian shape plus a polynomial. The $M_{\Xi^-\pi^-}$ distribution shown in Fig. 2 was fit (smooth curve) with this background form with a free normalization, together with a Gaussian shape whose position was fixed at 1862 MeV and whose width was fixed to the instrumental resolution of $\sigma = 10.2$ MeV, derived from a simulation of the spectrometer described below. The upper limit at the 90% confidence level (C.L.) for the hypothetical $\Xi_{3/2}^{--}$ peak is found to be 3.9 events. The $M_{\Xi^-\pi^+}$ distribution was fit (curve in Fig. 3) using two peak shapes plus the functional




FIG. 1 (color online). (a) Invariant mass distribution of the $p\pi^-$ system. (b) Invariant mass distribution of the $p\pi^-\pi^-$ system. Both invariant mass distributions contain also the charge conjugate (c.c.) modes ($\bar{p}\pi^+$, $\bar{p}\pi^+\pi^+$) and are subject to the constraints in event topology discussed in the text.




FIG. 2 (color online). Invariant mass distribution of the $p\pi^-\pi^-\pi^-$ system, subject to the constraints in event topology discussed in the text. The mixed-event background is represented by the gray shaded histogram, which is normalized to the background component of the fitted curve described in the text. The arrow shows the hypothetical $\Xi_{3/2}^{--}$ mass.

form that represents the mixed-event background. One peak shape was Gaussian with a position fixed at 1862 MeV and a fixed width of $\sigma = 10.2$ MeV, while the other was a convolution of Gaussian and Breit-Wigner shapes that was allowed to vary freely near the position of the well-established $\Xi^0(1530)$. This Gaussian contribution had a width fixed at 7.2 MeV, while the Breit-Wigner shape had the width fixed at the known $\Gamma = 9.1$ MeV for the $\Xi^0(1530)$ [32]. The position of the $\Xi^0(1530)$ at $1536.5 \pm 3.6(\text{stat})$ MeV is consistent with the expected value of 1531.8 MeV. The values for both Gaussian widths correspond to the simulated instrumental resolutions. The area of the $\Xi^0(1530)$ peak is found to be 35 ± 11 events. For the hypothetical $\Xi_{3/2}^0$ peak at 1862 MeV, the 90% C.L. upper limit is found to be 4.6 events. All of these results are from unbinned maximum likelihood fits [33] to the original event distributions.

In order to derive cross sections, the detector acceptance was simulated using two alternative models for the unknown production kinematics. In the first ansatz, the unknown kinematic distribution of the parent was described by functional forms resembling the transverse and longitudinal momentum distributions of Λ hyperons observed at HERMES. For the second model, the kinematic distributions of the parent were taken to be those of the $\Xi^0(1530)$




FIG. 3 (color online). Invariant mass distribution of the $p\pi^-\pi^-\pi^+$ (plus c.c.) system, subject to the constraints in event topology discussed in the text. The mixed-event background is represented by the gray shaded histogram, which is normalized to the background component of the fitted curve described in the text. The arrow shows the hypothetical $\Xi_{3/2}^0$ mass. The excess near 1.77 GeV has a statistical significance of only 1.8σ .

as predicted by the PYTHIA6 Monte Carlo code [34]. Both models were isotropic in the decay angle distribution.

In the case of the $\Xi^0(1530)$, the acceptance times efficiency from these two approaches are $0.036\%^1$ and 0.10% , respectively. This range in acceptance values due to the unknown production mechanism was carried forward into the cross section calculations. The photoproduction cross sections were evaluated using the Weizsäcker-Williams flux factor calculated over the range $3.6 < \nu < 24.6$ GeV. If the branching ratio of $\Xi^0(1530) \rightarrow \Xi^-\pi^+$ decay is taken to be $2/3$ [35], its photoproduction cross section is found to be between 8.8 and 24 nb. This experimental result is considerably higher than the PYTHIA6 model prediction [34] in 4π acceptance of 0.92 nb for production of $\Xi^0(1530)$ plus $\Xi^0(1530)$. However, the cross section ratio of $\Xi^0(1530)$ to $\Xi^0(1530)$ production predicted by PYTHIA6 is 1:4, which is consistent with the present data.

¹The acceptance for the 4-track decay of the $\Xi^0(1530)$ resonance is comparable to that for the 3-track decay of the Θ^+ [10] due to the inclusion of short tracks. These are tracks that make it through the spectrometer magnet, but not all the way to the RICH detector.

To simulate the detector response for the hypothetical $\Xi_{3/2}^{--}$ and $\Xi_{3/2}^0$ resonances, the particles were generated with a mass of 1862 MeV and an intrinsic width of $\Gamma = 2$ MeV, at vertices distributed according to the HERMES target gas profile. For the initial kinematic distributions, the same two models as for the $\Xi^0(1530)$ were employed. The instrumental resolution derived from this simulation was $\sigma = 10.2$ MeV in the $M_{\Xi^-\pi^-}$ and $M_{\Xi^-\pi^+}$ mass spectra near 1860 MeV.

Estimates of the spectrometer acceptance times efficiency from the simulation were used to estimate an upper limit for the cross section for production of the Ξ^{--} and Ξ^0 . Employing the two models described above for the initial kinematic distributions, the acceptances for these particles were estimated to be 0.031% and 0.065%, respectively. Using the fit results described above, the corresponding results for the upper limit for the Ξ^{--} (Ξ^0) photoproduction cross section times branching fraction is found to be 1.0 to 2.1 nb (1.2 to 2.5 nb) at the 90% C.L.. Based on the cross section ranges for the $\Xi^0(1530)$ and Ξ^{--} reported here, a 90% C.L. upper limit on the cross section ratio of $\Xi^{--} \rightarrow \Xi^-\pi^-$ to $\Xi^0(1530)$ between 0.06 and 0.15 has been derived. The values were obtained by assuming a central value for each cross section range, assigning the uncertainty to be half of the range, and performing error propagation on the ratio.

In summary, a search for the exotic Ξ^{--} baryon resonance has been performed in quasireal photoproduction on a deuterium target through the decay channel $\Xi^-\pi^- \rightarrow \Lambda\pi^-\pi^- \rightarrow p\pi^-\pi^-\pi^-$. The upper limit for the Ξ^{--} photoproduction cross section is 2.1 nb (90% C.L.). In addition, the well-established $\Xi^0(1530)$ is clearly identified in this experiment, and its photoproduction cross section is found to be between 8.8 and 24 nb, the range being due to the effect of the uncertainty in the production kinematics on the acceptance correction.

We gratefully acknowledge the DESY management for its support and the staff at DESY and the collaborating institutions for their significant effort. This work was supported by the FWO-Flanders, Belgium; the Natural Sciences and Engineering Research Council of Canada; the National Natural Science Foundation of China; the INTAS and ESOP network contributions from the European Community; the German Bundesministerium für Bildung und Forschung (BMBF); the Deutsche Forschungsgemeinschaft (DFG); the Deutscher Akademischer Austauschdienst (DAAD); the Italian Istituto Nazionale di Fisica Nucleare (INFN); Monbusho International Scientific Research Program, JSPS, and Toray Science Foundation of Japan; the Dutch Foundation for Fundamenteel Onderzoek der Materie (FOM); the U.K. Engineering and Physical Sciences Research Council and the Particle Physics and Astronomy Research Council; and the U.S. Department of Energy and the National Science Foundation.

- [1] D. Diakonov, V. Petrov, and M. Polyakov, *Z. Phys. A* **359**, 305 (1997).
- [2] M. Chemtob, *Nucl. Phys.* **B256**, 600 (1985).
- [3] H. Walliser, *Nucl. Phys.* **A548**, 649 (1992).
- [4] LEPs Collaboration, T. Nakano *et al.*, *Phys. Rev. Lett.* **91**, 012002 (2003).
- [5] DIANA Collaboration, V. V. Barmin *et al.*, *Phys. At. Nucl.* **66**, 1715 (2003); *Yad. Fiz.* **66**, 1763 (2003).
- [6] CLAS Collaboration, S. Stepanyan *et al.*, *Phys. Rev. Lett.* **91**, 252001 (2003).
- [7] SAPHIR Collaboration, J. Barth *et al.*, *Phys. Lett. B* **572**, 127 (2003).
- [8] A. E. Asratyan, A. G. Dolgolenko, and M. A. Kubantsev, *Phys. At. Nucl.* **67**, 682 (2004); *Yad. Fiz.* **67**, 704 (2004).
- [9] CLAS Collaboration, V. Kubarovsky *et al.*, *Phys. Rev. Lett.* **92**, 032001 (2004); **92**, 049902(E) (2004).
- [10] HERMES Collaboration, A. Airapetian *et al.*, *Phys. Lett. B* **585**, 213 (2004).
- [11] ZEUS Collaboration, S. Chekanov *et al.*, *Phys. Lett. B* **591**, 7 (2004).
- [12] SVD Collaboration, A. Aleev *et al.*, hep-ex/0401024.
- [13] COSY-TOF Collaboration, M. Abdel-Bary *et al.*, *Phys. Lett. B* **595**, 127 (2004).
- [14] P. Zh. Aslanyan, V. N. Emelyanenko, and G. G. Rikhhkvizkaya, hep-ex/0403044.
- [15] Yu. A. Troyan *et al.*, hep-ex/0404003.
- [16] A. R. Dzierba *et al.*, *Phys. Rev. D* **69**, 051901 (2004).
- [17] *Babar* Collaboration, B. Aubert *et al.*, hep-ex/0408064; FOCUS Collaboration, K. Stenson *et al.*, hep-ex/0412021.
- [18] BES Collaboration, J. Z. Bai *et al.*, *Phys. Rev. D* **70**, 012004 (2004).
- [19] PHENIX Collaboration, C. Pinkenburg *et al.*, *J. Phys. G* **30**, S1201 (2004).
- [20] HERA-B Collaboration, K. T. Knöpfle *et al.*, *J. Phys. G* **30**, S1363 (2004).
- [21] SPHINX Collaboration, Yu. M. Antipov *et al.*, *Eur. Phys. J. A* **21**, 455 (2004).
- [22] ALEPH Collaboration, S. Schael *et al.*, *Phys. Lett. B* **599**, 1 (2004).
- [23] HyperCP Collaboration, M. J. Longo *et al.*, *Phys. Rev. D* **70**, 111101 (2004).
- [24] NA49 Collaboration, C. Alt *et al.*, *Phys. Rev. Lett.* **92**, 042003 (2004).
- [25] R. Jaffe and F. Wilczek, *Phys. Rev. Lett.* **91**, 232003 (2003).
- [26] H. G. Fischer and S. Wenig, *Eur. Phys. J. C* **37**, 133 (2004).
- [27] WA89 Collaboration, M. I. Adamovich *et al.*, *Phys. Rev. C* **70**, 022201(R) (2004).
- [28] HERA-B Collaboration, I. Abt *et al.*, *Phys. Rev. Lett.* **93**, 212003 (2004).
- [29] ZEUS Collaboration, S. V. Chekanov *et al.*, hep-ex/0405013; CDF Collaboration, I. Gorelov *et al.*, hep-ex/0408025; E690 Collaboration, D. Christian *et al.*, Quarks and Nuclear Physics 2004, Bloomington, Indiana, 2004, <http://www.qnp2004.org/>; FOCUS Collaboration, K. Stenson *et al.*, hep-ex/0412021.
- [30] HERMES Collaboration, K. Ackerstaff *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **417**, 230 (1998).
- [31] N. Akopov *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **479**, 511 (2002).
- [32] Particle Data Group, S. Eidelman *et al.*, *Phys. Lett. B* **592**, 1 (2004).
- [33] W. Verkerke and D. Kirkby, physics/0306116.
- [34] T. Sjöstrand *et al.*, *Comput. Phys. Commun.* **135**, 238 (2001).
- [35] WA89 Collaboration, M. I. Adamovich *et al.*, *Eur. Phys. J. C* **11**, 271 (1999).