

Nongeometric fluxes as supergravity backgrounds

Fernando Marchesano¹ and Waldemar Schulgin²

¹*ASC, Ludwig-Maximilians-Universität, Theresienstraße 37, 80333 München, Germany*

²*Max Planck Institut für Physik, Föhringer Ring 6, 80805 München, Germany*

(Received 10 May 2007; published 30 August 2007)

We consider examples of $D = 4$ string theory vacua which, although globally nongeometric, admit a local description in terms of $D = 10$ supergravity backgrounds. We analyze such backgrounds and find that the supersymmetry spinors vary nontrivially along the internal manifold, reproducing the interpolating supergravity solutions found by Frey and Graña. Finally, we propose a simple, local expression for nongeometric fluxes in terms of the internal spinors of the compactification.

DOI: [10.1103/PhysRevD.76.041901](https://doi.org/10.1103/PhysRevD.76.041901)

PACS numbers: 11.25.Mj, 04.65.+e

Undoubtedly, fluxes have played a major role in recent attempts to connect string theory to observable physics. Why this is the case may be understood from two simple facts. First, string compactifications with background fluxes enlarge nontrivially the set of $D = 4$ string vacua based on conventional Calabi-Yau (CY) compactifications, and so the set of effective field theories that one can obtain. Second, flux compactifications are rich sources of $D = 4$ effective superpotentials, allowing to address two important issues of string phenomenology, moduli stabilization, and supersymmetry breaking, in a controlled manner [1].

A celebrated class of flux vacua is given by type IIB string theory compactified on a warped Calabi-Yau manifold with orientifold planes [2,3]. In many senses, one can treat the resulting $D = 4$ effective theory as the fluxless Calabi-Yau compactification, to which we add a flux-induced superpotential [4] and a nontrivial warping. This provides a simple global picture of this class of vacua, which even allows to treat them as an ensemble [5]. Finally, a nice feature of this set of $D = 4$ vacua is that they can be understood from $D = 10$ supergravity. The geometric intuition that one obtains from this fact has triggered several phenomenologically interesting scenarios, in particular, some related to early universe cosmology [6].

While warped CY manifolds have received a lot of attention, they are not the only class of flux compactifications in the literature, even if we restrict to type IIB $\mathcal{N} = 1$ $D = 4$ Poincaré invariant vacua. Another well-known family is based on compact non-Kähler manifolds, dual to the solutions found in the heterotic string [7]. Here the fluxes not only consist of field strengths, but also of torsion factors that deviate the metric from a Calabi-Yau one, and so they are usually referred to as geometric fluxes [8,9].

While geometric fluxes are less intuitive than the usual ones, an even more exotic kind of fluxes are those named nongeometric fluxes, which arise in the context of nongeometric compactifications [10–14]. Such compactifications are possible because, as string theory has a larger duality group than plain supergravity, it can be compactified in spaces where (classical) gravity would not even make sense. Perhaps the best understood class of nongeometric

metric vacua are those based on T folds [11], where locally there is a standard space-time description of the compactification but, as we relate two overlapping patches, we must use transition functions that mix the metric with the antisymmetric B field and interchange Kaluza-Klein (KK) with winding modes, just as T dualities do.

Clearly, nongeometric compactifications are a novel and exciting area to develop string phenomenology, as they are genuine string compactifications that cannot be described as $D = 10$ supergravity plus localized sources. Moreover, as nongeometric fluxes are combined with the previous fluxes, the effective superpotential that one obtains generalizes that of a Calabi-Yau manifold with fluxes [12], in such a way that compactifications with all moduli stabilized are easily obtained [15].

On the other hand, our current understanding of nongeometric compactifications is still too weak in order to perform a detailed analysis of their phenomenological possibilities. Part of the problem is that, unlike for other fluxes, there is no simple, intuitive definition of what nongeometric fluxes mean. In practice, they appear as the structure constants of $D = 4$ effective gauged supergravities, or as internal monodromies between the metric and B field [9,12,13]. In fact, very few examples of nongeometric compactifications have been built beyond Scherk-Schwarz compactifications and close relatives and, because of the issues above, it is hard to figure out how the space of nongeometric vacua looks.

The purpose of this paper is to shed further light into our understanding of nongeometric fluxes by describing them in terms of supergravity backgrounds. More precisely, we focus on some simple examples of T folds, where a local $D = 10$ description is possible, and study in detail the supergravity solutions associated to them. This will not only provide us with a simple definition of nongeometric flux, but will also unveil many local properties of this kind of backgrounds. Notice that the latter point is essential in order to build realistic scenarios from nongeometric compactifications. Indeed, most of the cosmological models constructed from warped CY's with fluxes are based on D-brane inflation, where only a local description of the su-

pergravity background (like, e.g., the Klebanov-Strassler (KS) throat [16]) is needed. Hence, based on the results below, similar scenarios could presumably be constructed for nongeometric vacua.

In order to perform our analysis, let us first construct a nongeometric vacuum. As pointed out in [9] this can be done by performing appropriate T dualities on simple geometric flux compactifications. In particular, let us consider type IIB string theory compactified on the warped toroidal background [17]

$$ds^2 = Z^{-1/2} ds_{M_4}^2 + Z^{1/2} ds_{T^6}^2, \quad (1)$$

$$ds_{T^6}^2 = \sum_{i=1}^3 ds_{(T^2)_i}^2, \quad ds_{(T^2)_i}^2 = R_i^2 |dx^i + \tau^i dx^{i+3}|^2, \quad (2)$$

$$H_3 = N(dx^1 \wedge dx^5 - dx^4 \wedge dx^2) \wedge dx^6, \quad (3)$$

$$F_3 = N(dx^4 \wedge dx^2 - dx^1 \wedge dx^5) \wedge dx^3, \quad (4)$$

$$\tilde{F}_5 = (1 + *_{10}) d\text{vol}_{M_4} \wedge dh, \quad (5)$$

$$\tau = C_0 + ie^{-\phi_0} = \text{const.}, \quad (6)$$

which is closely related to the vacua considered in [9], and a simple example of a warped Calabi-Yau manifold with H_3 and F_3 fluxes of [3]. Here $ds_{M_4}^2$ stands for the $D = 4$ Minkowski metric, τ for the type IIB axio-dilaton, the warp factor Z only depends on the internal coordinates $x^i \sim x^i + 1$, N is an even integer [18], and we must impose $dh = e^{-\phi_0} dZ^{-1}$ [3]. In order to find a solution to the Bianchi identity of $\tilde{F}_5$ and Einstein's equations we need to add negative tension objects to the compactification [3], which in our case will be 64 O3 planes filling up M_4 . Finally, by adding $32 - 2N^2$ D3 branes and no further localized objects all consistency conditions will be satisfied.

The presence of the background fluxes H_3 and F_3 generates a superpotential [4]

$$W = \int_{T^6} (F_3 - \tau H_3) \wedge \Omega_3 = N(\tau^1 - \tau^2)(\tau^3 - \tau) \quad (7)$$

and so we also need to impose $\tau^1 = \tau^2$ and $\tau^3 = \tau$. This will guarantee that the complexified 3-form flux $G_3 = F_3 - \tau H_3$ is an imaginary-self-dual (ISD), primitive (2, 1)-form, and so we will have a supersymmetric vacuum [19]. In fact, due to the simplicity of our example, the present compactification yields a $D = 4$ $\mathcal{N} = 2$ effective theory. The supersymmetry generators are of the form

$$\epsilon_L = \frac{1}{\sqrt{2}}(u \otimes \chi_L + u^* \otimes \chi_L^*), \quad (8)$$

$$\epsilon_R = \frac{1}{\sqrt{2}}(u \otimes \chi_R + u^* \otimes \chi_R^*), \quad (9)$$

where u stands for the external and χ for the internal space-time spinor, the latter chosen of negative chirality. While the above *Ansatz* is valid for general supersymmetric type IIB compactifications, here the presence of the flux G_3 and the O3 planes imposes the following relation

$$\epsilon_L = \Gamma_{O3} \epsilon_R = -i\Gamma^{(6)} \epsilon_R \quad (10)$$

between left (ϵ_L) and right (ϵ_R) spinors. Here $\Gamma^{(6)}$ stands for the internal chirality operator, and so we have that

$$\chi_L = i\chi_R \rightarrow \epsilon = \epsilon_L + i\epsilon_R = \sqrt{2}u \otimes \chi_L \quad (11)$$

which is the usual B-type *Ansatz* for warped Calabi-Yau compactifications with fluxes [20]. In the present background, the internal spinors are of the form $\chi_{R,i} = Z^{-1/8} \eta_i$, where

$$\eta_1 = \frac{1}{2}(- - -), \quad \eta_2 = \frac{1}{2}(+ + -) \quad (12)$$

in the usual spinor notation.

As pointed out in [9], by performing two consecutive T dualities along two directions of an H_3 component one obtains a nongeometric compactification. In particular, following the notation of [12], after 2 T dualities along $\{x^5, x^1\}$ the flux $H_3 = dB$ partially becomes a nongeometric Q flux via

$$H_{x^1 x^5 x^6} = N \xrightarrow{T_{x^5,1}} Q_{x^6}^{x^5 x^1} = N, \quad (13)$$

whereas the other component of H_3 remains untouched.

In order to obtain the new background one can apply Buscher's T-duality rules [21], but first one needs to have the directions $\{x^5, x^1\}$ as isometries of the background. This is achieved by choosing an appropriate gauge for the B field, like $B = Nx^6(dx^1 \wedge dx^5 - dx^4 \wedge dx^2)$, and by smoothing out the localized sources (i.e., the D3 branes and O3 planes) along $\{x^5, x^1\}$. This means, in particular, that the warp factor Z will be given by a function $\Delta \equiv \Delta(x^2, x^3, x^4, x^6)$, which by Einstein's equations must satisfy

$$-e^{-\phi_0} \nabla_{T^4}^2 \Delta = \frac{2N^2}{\text{Vol}_{T^4}} + \sum_i q_i \delta_{T^4}(\vec{x}_i), \quad (14)$$

where T^4 stands for a four-torus of coordinates $\{x^2, x^4, x^3, x^6\}$ and the delta functions represent the sources localized in the same T^4 , with a charge $q = 1$ in the case of a D3 brane and $q = -2$ for a (partially delocalized) O3 plane. Notice that on T^4 the O3 planes are at the fixed locations $x^i \in \mathbb{Z}/2$ ($i = 2, 3, 4, 6$), and in the following we will focus our attention to the neighborhood containing the one at the origin. Finally, in order to simplify the discussion below, we will choose the compactification moduli to be $R_1 = R_2 = 1$ and $\tau^1 = \tau^2 = i$, although more general choices consistent with the supersymmetry condition $\tau^1 = \tau^2$ may also be chosen.

Taking into account these prescriptions and directly applying Buscher's rules along $\{x^5, x^1\}$ one again obtains type IIB string theory, but this time on the background

$$ds^2 = \Delta^{-1/2} ds_{M_4}^2 + \Delta^{1/2} ds_{M_6}^2, \quad (15)$$

$$ds_{M_6}^2 = (dx^2)^2 + (dx^4)^2 + d\bar{z}^3 d\bar{z}^3 + e^{2(\phi - \phi_0)} ((dx^1)^2 + (dx^5)^2), \quad (16)$$

$$B = Nx^6(dx^2 \wedge dx^4 + e^{2(\phi-\phi_0)}dx^5 \wedge dx^1), \quad (17)$$

$$F_1 = N \text{Red} z^3, \quad (18)$$

$$\tilde{F}_3 = F_1 \wedge B + e^{2\phi-\phi_0} *_{\mathcal{M}_6} de^{-2\phi} \wedge dx^5 \wedge dx^1, \quad (19)$$

$$\tilde{F}_5 = (1 + *_{10})d\text{vol}_{M_4} \wedge d(Nx^6/\Delta)e^{-\phi_0}, \quad (20)$$

$$e^{\phi-\phi_0} = (\Delta + (Nx^6)^2)^{-1/2}, \quad (21)$$

where for simplicity we have defined $dz^3 = dx^3 + \tau dx^6$, with τ given by (6). When compactifying this theory, one should perform the identifications $x^3 \sim x^3 + 1$ and $x^6 \sim x^6 + 1$. However, for reasons that will become clear below, we will initially not impose such identifications. On the other hand, we will still impose $x^i \sim x^i + 1$ for $i = 1, 2, 4, 5$. The background fluxes $\tilde{F}_3 = F_3 - C_0 H_3$, $F_1 = dC_0$ stand for the generalized field strength of type IIB supergravity. Notice that neither the axion C_0 nor the dilaton $e^{-\phi}$ are constant in this new background. Nevertheless, as usual one can still define an average string coupling by setting $g_s = e^{\phi_0}$, with e^{ϕ_0} given by (6), that can be taken to arbitrary small values.

Just as the above background has been T dualized, we must also transform the localized sources, namely, the orientifold planes and the open string sector of the theory. By the usual T-duality rules we would now expect to have O5 planes and D5 branes wrapped on the two-torus spanned by $\{x^5, x^1\}$. This intuition is confirmed by looking at the Bianchi identities of the new background. While $dF_1 = dH_3 = 0$ are clearly satisfied, for $\tilde{F}_3$ we have that [22]

$$\begin{aligned} d\tilde{F}_3 - H_3 \wedge F_1 &= g_s^{-1} \nabla_{\mathbf{T}^4}^2 e^{-2(\phi-\phi_0)} d\text{vol}_{\mathbf{T}^4} \\ &= -\sum_i q_i \delta_{\mathbf{T}^4}(x_i) d\text{vol}_{\mathbf{T}^4}, \end{aligned} \quad (22)$$

where we have used (14) and (19), and we are defining $d\text{vol}_{\mathbf{T}^4} = R_3^2 \text{Im}\tau dx^2 \wedge dx^4 \wedge dx^3 \wedge dx^6$. Hence we have a set of localized sources to which the Ramond-Ramond (RR) field strength F_3 couples. As we expected, these are D5 branes and O5 planes, wrapping $M_4 \times \mathbf{T}^2_{(x^5, x^1)}$ and being pointlike in $\{x^2, x^4, x^3, x^6\}$. In particular, there will be an O5 plane at the origin $x^2 = x^3 = x^4 = x^6 = 0$ and, if we treat such coordinates as noncompact, one at each location given by $x^i \in \mathbb{Z}/2$ ($i = 2, 3, 4, 6$), as we would also expect from T-duality. The remaining Bianchi identity can be easily computed if we realize that the internal component of $\tilde{F}_5$ satisfies

$$\tilde{F}_5 - B \wedge \tilde{F}_3 + \frac{1}{2} B^2 \wedge F_1 = -e^{-2(\phi-\phi_0)} N *_{\mathcal{M}_6} \text{Im} dz^3 \quad (23)$$

and so the right-hand side (r.h.s.) of (23) is closed. This implies that

$$d\tilde{F}_5 - H_3 \wedge \tilde{F}_3 = B \wedge (d\tilde{F}_3 - H_3 \wedge F_1) \quad (24)$$

and so, by (22), we again have the sum of delta functions.

The presence of the B field is easy to understand from the fact that the left-hand side (l.h.s.) of (24) measures the D3-brane charge of localized sources, and this same charge is induced on D5 branes in the presence of a nontrivial B field [23]. Hence the r.h.s. of (24) takes into account that we are in the presence of D5 branes magnetized by a nontrivial B field. What is perhaps more surprising is the fact that the O5 planes which are not located at $x^6 = 0$ also seem to be “magnetized.” In that case we should be dealing with exotic orientifold planes, analogous to the ones analyzed in [24]. Finally, in the same way that we have solved the Bianchi identities, one can check that the equations of motion for the metric, axio-dilaton, and background fluxes are satisfied. As a result, even if one is skeptical about applying Buscher’s rules on the initial type IIB background, it is clear that Eqs. (15)–(21) describe a genuine type IIB supergravity vacuum.

Notice that in this new background the warp factor Z' and the 4-form potential $C'_4 = h' d\text{vol}_{M_4}$ are given by

$$Z' = \Delta, \quad h' = Nx^6 e^{-\phi_0} \Delta^{-1} \quad (25)$$

and so the usual relation $d(h - e^{-\phi} Z^{-1}) = 0$ of type IIB compactifications with ISD G_3 fluxes, D3 branes, and O3 planes is no longer satisfied. Such a relation is only satisfied in the limit $x^6 \rightarrow \infty$, where the B field is so strong and the two-torus $\{x^5, x^1\}$ is so small that a D5 brane wrapped on it looks like a D3 brane. On the other hand, on the limit $x^6 \rightarrow -\infty$ we recover the relation $d(h + e^{-\phi} Z^{-1}) = 0$, satisfied by type IIB backgrounds with IASD G_3 fluxes, anti-D3 branes and anti-O3 planes. Finally, at $x^6 = 0$ we have that $h = 0$ and $Z = e^{-2(\phi-\phi_0)}$, which is typical of type IIB flux compactifications containing O5 planes and geometric fluxes.

It thus seems that, as we move along x^6 , our type IIB background interpolates between different classes of type IIB compactifications. We can parametrize such an interpolation by defining an angle α as

$$\sin \alpha = \Delta^{1/2} e^{\phi-\phi_0}, \quad \cos \alpha = Nx^6 e^{\phi-\phi_0}, \quad (26)$$

and analyzing how the background depends on it. A non-trivial dependence will appear on the forms J and Ω describing the metric, which can be taken to be

$$\begin{aligned} \frac{J}{\Delta^{1/2}} &= e^{2(\phi-\phi_0)} dx^5 \wedge dx^1 + dx^2 \wedge dx^4 + i \frac{R_3^2}{2} dz^3 \wedge d\bar{z}^3, \\ \frac{\Omega}{\Delta^{3/4}} &= i e^{\phi-\phi_0} R_3 (dx^1 + i dx^5) \wedge (dx^4 + i dx^2) \wedge dz^3, \end{aligned}$$

which naturally define a complex and a symplectic structure. In terms of these definitions, one can check that the ISD, (2, 1)-form in this background is

$$\mathcal{G}_3 = \tilde{F}_3 - i e^{-\phi} \text{Re}(e^{i\alpha} [H_3 - i \Delta^{1/2} d(\Delta^{-1/2} J)]), \quad (27)$$

which, at specific values of α takes a more familiar form

$$\alpha \rightarrow 0 \Rightarrow \mathcal{G}_3 = \tilde{F}_3 - i e^{-\phi} H_3 = G_3, \quad (28)$$

$$\alpha = \pi/2 \Rightarrow \mathcal{G}_3 = \tilde{F}_3 - ie^{-2\phi} d(e^\phi J), \quad (29)$$

$$\alpha \rightarrow \pi \Rightarrow \mathcal{G}_3 = \tilde{F}_3 + ie^{-\phi} H_3 = \tilde{G}_3. \quad (30)$$

Finally, a D5 brane wrapped on $\{x^5, x^1\}$ satisfies

$$\iota_X \Omega|_{D5} = \text{Im}(e^{-i\alpha}(B + iJ))|_{D5} = 0, \quad (31)$$

which are the usual Bogomol'nyi-Prasad-Sommerfield (BPS) conditions for a D5 brane in a Calabi-Yau compactification [25]; notice, however, that in the Calabi-Yau case α is a constant parameter, while here it varies along the compactification manifold.

But the easiest way to see that this supergravity background interpolates between different classes of standard type IIB compactifications is by analyzing the supersymmetry spinors. These spinors can also be obtained by T duality, by applying the rules of [26]. One then obtains that the new spinors no longer satisfy the relation (11), but instead

$$\epsilon'_L = (\cos\alpha\Gamma_{03} + \sin\alpha\Gamma_{05})\epsilon'_R, \quad (32)$$

where Γ_{05} is the chirality operator on the coordinates $\{x^2, x^4, x^3, x^6\}$ transverse to $M_4 \times \mathbf{T}^2_{(x^5, x^1)}$, as well as the orientifold projection that an O5 plane wrapping $M_4 \times \{x^5, x^1\}$ would implement in standard compactifications.

Now, since our initial background contained two independent spinors, so will the new one. In order to see how the $D = 10$ spinors (8) and (9) now look, let us consider a particular linear combination of them. Namely, we choose

$$\chi_{R,\pm} = \Delta^{-1/8} \eta_{\pm}, \quad \eta_{\pm} = \frac{1}{\sqrt{2}}(\eta_1 \pm \eta_2) \quad (33)$$

then, taking into account that

$$\Gamma_{03}\eta_{\pm} = i\eta_{\pm}, \quad \Gamma_{03}\eta_{\pm}^* = -i\eta_{\pm}^*, \quad (34)$$

$$\Gamma_{05}\eta_{\pm} = \pm\eta_{\pm}, \quad \Gamma_{05}\eta_{\pm}^* = \pm\eta_{\pm}^*, \quad (35)$$

we obtain the following $D = 10$ spinors

$$\epsilon_{R,\pm} = \frac{1}{\sqrt{2}}(u \otimes \chi_{R,\pm} + u^* \otimes \chi_{R,\pm}^*), \quad (36)$$

$$\epsilon_{L,\pm} = \frac{1}{\sqrt{2}}(ie^{\pm i\alpha} u \otimes \chi_{R,\pm} - ie^{\mp i\alpha} u^* \otimes \chi_{R,\pm}^*) \quad (37)$$

precisely matching the spinor *Ansatz* considered in [27], where type IIB supergravity solutions interpolating between standard compactification *Ansätze* were shown to be possible.

In particular, the interpolating solutions analyzed in [27] connected backgrounds where D3 branes are BPS objects to those where D5 branes are BPS. We see that, if we extend our supergravity solution along $x^6 \in (-\infty, \infty)$ the same situation will happen, with the difference that now anti-D3 branes will also be included. So, just as expected, our supergravity background is such that for $x^6 \rightarrow -\infty$ anti-D3 branes are BPS, for $x^6 = 0$ D5 branes are BPS, and for $x^6 \rightarrow \infty$ D3 branes are BPS. At intermediate points

of x^6 the supersymmetry preserved by the background is that of a magnetized D5 brane.

This description fits exactly with the D5-brane BPS conditions found in (31), which can be derived as follows. First, one follows the computations of [27] in order to see that (36) and (37) are indeed supersymmetry generators of the new background. Second, one chooses one of the two spinors, say η_+ , and constructs the spinor bilinears

$$J_{mn} = -i\eta_+^\dagger \Gamma_{mn} \eta_+, \quad \Omega_{mnp} = -\eta_+^T \Gamma_{mnp} \eta_+ \quad (38)$$

familiar from $SU(3)$ -structure compactifications [1], and obtains the same forms J and Ω as above. Finally, by repeating the κ -symmetry computations of [25], one arrives at (31). Had we chosen the internal spinor η_- , we would have obtained a different 2-form J , but nevertheless the same final condition for a D5 brane wrapped on $\{x^5, x^1\}$. Alternatively, the same conclusion can be reached by using the results in [28].

To sum up, we have analyzed the type IIB supergravity backgrounds associated to a simple set of nongeometric backgrounds, and found that they correspond to solutions that interpolate between standard type IIB compactifications. This interpolation is particularly manifest in the internal part of the space-time supersymmetry generators, which fit into the *Ansatz* used in [27]. Although our background is particularly simple and yields $D = 4$ extended supersymmetry, we do not expect this to be the general case. In fact, by analyzing other simple nongeometric backgrounds we find that the *Ansatz* (36) and (37) is not always realized, but rather more general interpolating *Ansätze* like that used in [29] and generalizations. What remains valid is the relation (32) between left and right-handed spinors, in the sense that we have a rotation between two usual orientifold projections Γ_{Op} and Γ_{Oq} with a rotation parameter α varying along the internal manifold. This should not only be true for toroidal-like nongeometric vacua, but also for more general ones that can be obtained from fiberwise T duality on a geometric flux background. It would be interesting to verify this by generalizing the computations performed in [30]. Remarkably, the rotation *Ansatz* (32) is compatible with the presence of orientifold planes, as our example explicitly shows. Indeed, by construction we know that there is an O5 plane located at $x^6 = 0$. At this point $\alpha = \pi/2$ and so (32) reduces to the usual O5-plane projection (in flat space), but notice that this is no longer true as soon as $x^6 \neq 0$.

When describing this background as a nongeometric compactification, $\{x^2, x^4, x^3, x^6\}$ parametrize a four-torus $\mathbf{T}^4$, over which the $\mathbf{T}^2$ with coordinates $\{x^5, x^1\}$ is fibered. As we perform the identification $x^6 \sim x^6 + 1$, $\mathbf{T}^2_{(x^5, x^1)}$ will suffer a nongeometric monodromy (namely a T-duality transformation) that signals the presence of the nongeometric Q flux (13). Notice that this is a global definition, and that there is no obvious local definition of a Q flux, like the one for a 3-form flux H_3 via $H_3 = dB$. However, our background example above suggests how such a definition

could be made. Consider the quantity

$$q_a^{bc} = \partial_a \left(\frac{\chi_R^\dagger \gamma^{bc} \Gamma_{Op} \chi_L}{e^{\phi - \phi_0} \chi_L^\dagger \chi_L} \right) \quad (39)$$

defined in a local, geometric neighborhood containing an Op plane, where γ^{bc} is the antisymmetrized product of two γ matrices. The usual Q flux is then obtained by integrating (39) over a one-cycle γ_a :

$$Q_a^{bc} = \int_{\gamma_a} q_a^{bc} \quad (40)$$

(see [31] for alternative, possibly related definitions). It is easy to see that, by plugging (26) and (32) in the above expressions and setting $p = 5$ one recovers the Q-flux component (13) present in our background.

It would be interesting to compute (39) and (40) for more involved nongeometric compactifications, as well as to generalize the notion of Q charge, along the lines of [32–34]. In any case, it should be clear what the intuition behind the above definitions is. The local quantity (39) measures

how the relative rotation generated by γ^{bc} between the left and right spinors varies along the internal manifold. Hence it will always vanish for the standard, geometric *Ansätze* [7,20]. On the other hand, if (40) does not vanish for some closed path γ_a , then there is a spin monodromy that acts differently for the internal spinors χ_L and χ_R . This could not possibly be for a geometric compactification, since in that case both χ_L and χ_R are the same kind of objects, i.e., sections of the same spin bundle, and should transform in the same way when going around γ_a . Thus, (39) and (40) provide a suitable way to measure global nongeometrical aspects of string compactifications. The hope is that, with these definitions at hand, one can better understand what the space of nongeometric vacua is.

We wish to thank A. Font, M. Graña, L. E. Ibáñez, and P. Koerber for useful discussions. F.M. is supported by the European Network “Constituents, Fundamental Forces and Symmetries of the Universe,” under the Contract No. MRTN-CT-2004-005104. W.S. would like to thank the University of Munich for hospitality.

-
- [1] For reviews see, e.g., M. Graña, Phys. Rep. **423**, 91 (2006); M.R. Douglas and S. Kachru, Rev. Mod. Phys. **79**, 733 (2007); R. Blumenhagen, B. Körs, D. Lüst, and S. Stieberger, arXiv:hep-th/0610327.
 - [2] K. Dasgupta, G. Rajesh, and S. Sethi, J. High Energy Phys. **08** (1999) 023.
 - [3] S.B. Giddings, S. Kachru, and J. Polchinski, Phys. Rev. D **66**, 106006 (2002).
 - [4] S. Gukov, C. Vafa, and E. Witten, Nucl. Phys. **B584**, 69 (2000); **B608**, 477(E) (2001).
 - [5] F. Denef, M.R. Douglas, and S. Kachru, arXiv:hep-th/0701050.
 - [6] For reviews see C. P. Burgess, arXiv:hep-th/0606020; S. H. Henry Tye, arXiv:hep-th/0610221; J. M. Cline, arXiv:hep-th/0612129; R. Kallosh, arXiv:hep-th/0702059.
 - [7] C. M. Hull, Print-86-0251-CAMBRIDGE; A. Strominger, Nucl. Phys. **B274**, 253 (1986).
 - [8] S. Gurrieri, J. Louis, A. Micu, and D. Waldram, Nucl. Phys. **B654**, 61 (2003).
 - [9] S. Kachru, M. B. Schulz, P. K. Tripathy, and S. P. Trivedi, J. High Energy Phys. **03** (2003) 061.
 - [10] A. Dabholkar and C. Hull, J. High Energy Phys. **09** (2003) 054; S. Hellerman, J. McGreevy, and B. Williams, J. High Energy Phys. **01** (2004) 024; A. Flournoy, B. Wecht, and B. Williams, Nucl. Phys. **B706**, 127 (2005).
 - [11] C. M. Hull, J. High Energy Phys. **10** (2005) 065.
 - [12] J. Shelton, W. Taylor, and B. Wecht, J. High Energy Phys. **10** (2005) 085.
 - [13] A. Dabholkar and C. Hull, J. High Energy Phys. **05** (2006) 009.
 - [14] G. Aldazábal, P. G. Cámara, A. Font, and L. E. Ibáñez, J. High Energy Phys. **05** (2006) 070.
 - [15] J. Shelton, W. Taylor, and B. Wecht, J. High Energy Phys. **02** (2007) 095; A. Micu, E. Palti, and G. Tasinato, J. High Energy Phys. **03** (2007) 104.
 - [16] I. R. Klebanov and M. J. Strassler, J. High Energy Phys. **08** (2000) 052.
 - [17] We are using the convention $4\pi^2\alpha' = 1$.
 - [18] A. R. Frey and J. Polchinski, Phys. Rev. D **65**, 126009 (2002).
 - [19] M. Graña and J. Polchinski, Phys. Rev. D **63**, 026001 (2000).
 - [20] K. Becker and M. Becker, Nucl. Phys. **B477**, 155 (1996).
 - [21] T. H. Buscher, Phys. Lett. B **194**, 59 (1987).
 - [22] Following the conventions in [3], for a p -form α on a manifold $\mathcal{M}$ we have that $\alpha \wedge *_{\mathcal{M}} \alpha = -\alpha \cdot \alpha \text{dvol}_{\mathcal{M}}$.
 - [23] M. R. Douglas, arXiv:hep-th/9512077.
 - [24] C. Angelantonj, E. Dudas, and J. Mourad, Nucl. Phys. **B637**, 59 (2002).
 - [25] M. Mariño, R. Minasian, G. W. Moore, and A. Strominger, J. High Energy Phys. **01** (2000) 005.
 - [26] S. F. Hassan, Nucl. Phys. **B568**, 145 (2000).
 - [27] A. R. Frey and M. Graña, Phys. Rev. D **68**, 106002 (2003).
 - [28] L. Martucci and P. Smyth, J. High Energy Phys. **11** (2005) 048.
 - [29] G. Dall’Agata, Nucl. Phys. **B695**, 243 (2004).
 - [30] S. Fianza, R. Minasian, and A. Tomasiello, Commun. Math. Phys. **254**, 401 (2005).
 - [31] I. Benmachiche and T. W. Grimm, Nucl. Phys. **B748**, 200 (2006); I. T. Ellwood, arXiv:hep-th/0612100.
 - [32] J. Gray and E. J. Hackett-Jones, J. High Energy Phys. **05** (2006) 071.
 - [33] M. Graña, J. Louis, and D. Waldram, J. High Energy Phys. **04** (2007) 101.
 - [34] R. D’Auria, S. Ferrara, and M. Trigiante, arXiv:hep-th/0701247.