

Investigation of resonances in gravity-capillary wave turbulence

Quentin Aubourg¹ and Nicolas Mordant^{1,2,*}

¹*Université Grenoble Alpes, LEGI, CNRS, F-38000 Grenoble, France*

²*Institut Universitaire de France, 103 Boulevard Saint Michel, F-75005 Paris, France*

(Received 13 January 2016; published 7 June 2016)

We report experimental results on nonlinear wave coupling in surface wave turbulence on water at scales close to the crossover between surface gravity waves and capillary waves. We study three-wave correlations either in the frequency domain or in the wave-vector domain. We observe that in a weakly nonlinear regime, the dominant nonlinear interactions correspond to waves that are collinear or close to collinear. Although the resonant coupling of pure gravity waves is supposed to involve four waves, at the capillary crossover we observe a nonlocal coupling between a gravity wave and two capillary waves. Furthermore, nonlinear spectral spreading permits three-gravity wave coupling. These observations raise the question of the relevance of these processes in the oceanographic context and in particular the range of frequencies of gravity waves that may be impacted.

DOI: [10.1103/PhysRevFluids.1.023701](https://doi.org/10.1103/PhysRevFluids.1.023701)

I. INTRODUCTION

The weak turbulence theory (WTT) has been developed since the 1960s with initial motivations in oceanography and plasmas physics (see [1–3] for reviews). It provides a path from the dynamical wave equations to a statistical theory and notably the time evolution of the wave spectrum. This major step toward a statistical theory of a turbulent system can only be achieved under the hypotheses of both asymptotically large systems and asymptotically weak nonlinearity. In this framework, a natural scale separation exists between the fast period of the oscillations of the waves and the slow nonlinear evolution due to coupling with the other waves. This separation permits one to develop an asymptotic statistical theory. Because of the weakness of the nonlinear coupling, only a resonant wave can develop a long time cumulative coupling and in this way exchange significant amounts of energy. These waves verify the resonance conditions on wave numbers $\mathbf{k}_i$ and angular frequencies ω_i ,

$$\mathbf{k}_1 = \mathbf{k}_2 + \mathbf{k}_3, \quad \omega_1 = \omega_2 + \omega_3, \quad (1)$$

in the case of three interacting waves. The number of interacting waves to consider is determined generally by the order of the nonlinear term (three-wave coupling for quadratic nonlinearity, four-wave coupling for cubic ones, etc.). Exceptions come from the curvature of the linear dispersion relation that sometimes prevents one from finding any solutions for the three-wave interaction, for instance, despite quadratic nonlinear terms. This is the case for dispersion relations $\omega \propto k^\alpha$ for $\alpha < 1$ as is the case for pure gravity surface waves for which $\omega = \sqrt{gk}$ (g being the acceleration of gravity). In such a case the four-wave resonance has to be investigated. In real systems that have a finite size, discretization of frequencies and wave vectors renders the situation even more complex as the resonant sets can become extremely restricted [4–7]. In real systems, the magnitude of the waves and thus of the nonlinear coupling is also finite. Energy exchanges between waves decrease the coherence of the waves and thus induce a finite spectral width of the modes. The resonances are thus approximate,

$$\mathbf{k}_1 \approx \mathbf{k}_2 + \mathbf{k}_3, \quad \omega_1 \approx \omega_2 + \omega_3, \quad (2)$$

as some uncertainty exists on the frequencies and the wave numbers due to nonlinear spectral widening. Depending on the amplitude of the mismatch, this set of approximate resonances is

*nicolas.mordant@ujf-grenoble.fr

possibly much larger than that of exact resonances and this may enable a much more efficient wave coupling and thus a much more efficient energy transfer among waves.

The notion of wave resonance is thus at the core of the WTT. Not only does it permit one to derive the evolution equations for statistical quantities such as spectra, but solutions of these equations can be exhibited in stationary situations notably in the forced and thus out-of-equilibrium case [1–3]. The phenomenology associated with the latter case is that of the energy cascade in scale from the forced scales (usually large scales) to the scales at which dissipation dominates and ultimately removes energy as heat. The stationary out-of-equilibrium spectrum has been called a Kolmogorov-Zakharov (KZ) spectrum in honor of Zakharov and because of the analogy with the Kolmogorov spectrum of Navier-Stokes turbulence. In the case of pure capillary waves (short-wavelength limit of surface waves on a liquid) this spectrum of the surface deformation can actually be predicted by dimensional analysis (and confirmed analytically) to be $E_\zeta(k) \propto \gamma^{1/4} P^{1/2} / k^{7/4}$, where $k = ||\mathbf{k}||$, γ is the surface tension, and P is the energy injected (and dissipated) in the stationary system. The exponent $1/2$ comes from the three-wave interaction. For pure gravity waves (long-wavelength limit) the spectrum is $E_\zeta(k) \propto g^{1/2} P^{1/3} / k^{5/2}$, the coefficient $1/3$ being due to four-wave resonances as discussed above. In the general case of capillary-gravity waves, i.e., for wavelengths of centimeter scale, the dispersion relation is

$$\omega = \sqrt{gk + \frac{\gamma}{\rho} k^3}. \quad (3)$$

As it is not a pure power law, the analytic derivation of the KZ spectrum is much more difficult. It has been shown [8,9] that for any given value of ω_2 three-wave resonances are possible but the interaction becomes quite nonlocal when it takes values in the gravity range. Preliminary examinations of resonances in the turbulent state have suggested that the interactions are much more efficient in the almost unidirectional situation in which all the wave vectors are close to being collinear [10].

In this article we investigate in more detail the wave coupling in the turbulent state. We investigate three-wave correlations in either the frequency or the wave-number domain. This statistical estimator permits one to check the existence of wave resonances. We study two regimes: a weakly nonlinear regime and a regime of stronger forcing. The experimental setup is described in Sec. II. General spectral characteristics of the wave field are presented in Sec. III. The resonances are specifically studied in Sec. IV: first the theoretical exact resonances and then correlations in the frequency domain or in wave-vector space. A summary and conclusions are given in Sec. V.

II. EXPERIMENTAL SETUP

A sketch of the global configuration is shown in Fig. 1. The experimental setup is made of a $57 \times 37 \text{ cm}^2$ wave tank filled with water at a rest height $h_0 = 5 \text{ cm}$. Surface waves are generated by an oscillating motion of the wave tank, achieved with an oscillating table. We drive the table with a constant amplitude oscillation, which is frequency modulated. The frequency explores the interval $[0.5, 1.5] \text{ Hz}$, which includes an eigenmode of the wave tank. Two convex walls are used to break the anisotropy of the forcing by divergent wave reflections. To perform a full measurement of waves in space and time, we use the Fourier transform profilometry [11]. A video projector projects a pattern on the water surface and a high-speed camera with a parallel optical axis is focused on the surface. For the image to form at the very surface, water must diffuse light. As suggested by Prządka *et al.* [12], we use anatase titanium dioxide particles (Kronos 1001). These particles do not cause alteration of the pure water surface tension. A concentration of about 40 g/l permits us to project a pattern at the water surface with a high contrast level. When waves are present, the pattern is deformed and recorded by the high-speed camera. The deformation of the pattern can then be inverted back to the deformation of the surface $\eta(x, y)$. Our measurement area covers a surface of $20 \times 20 \text{ cm}^2$ at the center of the tank, with 1024×1024 pixel resolution at 250 frames/s . Data sets are composed of 15 movies of 87 s each to get a good statistical convergence. In this way we obtain measurements of the surface deformation $\eta(x, y, t)$ that are resolved in time and space. We also study




FIG. 1. Sketch of the experimental setup. Waves are generated by horizontal oscillation of the vessel. Two convex walls are used to improve the isotropy by divergent wave reflection. The video projector creates a pattern at the surface, the deformation of which is recorded by a high-speed camera. The pattern can then be inverted to recover the deformation [11]. The measurement region is centered and covers a surface $20 \times 20 \text{ cm}^2$.

the normal velocity field $v(x, y, t) = \frac{\partial \eta}{\partial t}$. Fourier analysis can then be performed by applying a digital Fourier transform in time and/or space.

We show in Fig. 2 two snapshots of the vertical velocity field $v = \frac{\partial \eta}{\partial t}$, where $\eta(x, y, t)$ is the altitude of the water surface: (a) weak forcing and (b) stronger forcing. To get an estimation of the intensity of the forcing, we define the typical wave steepness σ :

$$\sigma = \left\langle \sqrt{\frac{1}{S} \int_S \|\nabla h(x, y, t)\|^2 dx dy} \right\rangle. \quad (4)$$




FIG. 2. (a) Vertical velocity field for an experiment for a weak forcing where the typical wave steepness is $\sigma = 0.025$. (b) Vertical velocity field for a stronger forcing where $\sigma = 0.08$.




FIG. 3. (a) Temporal spectrum $E^\eta(\omega) = \langle |\eta(x, y, \omega)|^2 \rangle$ for two experiments where the typical wave steepness is $\sigma = 0.025$ (bottom) and $\sigma = 0.08$ (top). Dashed red lines represent theoretical slopes predicted by the WTT for pure gravity waves (proportional to ω^{-4}) expected at low frequencies and pure capillary waves expected at large frequencies (proportional to $\omega^{-17/6}$). (b) Spectral exponent fitted for a set of experiments on the low-frequency part of our spectrum.

We observe that the stronger forcing is more organized with the presence of coherent waves packets. Our experiments range roughly from $\sigma = 0.02$ to $\sigma = 0.1$. The two regimes presented recover the two regimes of water turbulence described by Cobelli *et al.* [13]. At weak steepness, a regime of truly weakly nonlinear wave turbulence is observed, as will be confirmed below. At larger steepness a regime of strongly nonlinear waves is observed. Part of the strongly nonlinear structures that are observed could actually come directly from the forcing or interactions with the wall. Indeed, at very weak forcing, the amplitude of the waves is very small (a couple of millimeters), so the perturbations induced by the depinning of the water contact line on the asperities of the walls are no longer fully negligible. A strong wave amplitude can be generated as well by focusing in the corners of the box. At very strong forcing the velocity of the walls can become close to the celerity of the waves, so the waves that are generated can be strongly nonlinear at the generation. Thus separating the origin of the nonlinearity between bulk and walls is actually challenging, although one expects the bulk effects to dominate when the system is large enough.

III. SPECTRAL ANALYSIS

In this section we analyze the turbulence regime by using Fourier analysis. By computing the Fourier transform in a time of $v(x, y, t)$ we obtain the temporal spectrum $E^\eta(\omega) = \langle |\eta(x, y, \omega)|^2 \rangle$, where $\langle \dots \rangle$ denotes the average over time windows and space. We compute the time Fourier transform using the Welch method with a 16-s time window of the signal. Figure 3(a) shows $E^\eta(\omega)$ for the two experiments where $\sigma = 0.025$ and $\sigma = 0.08$. The forcing operates in the range [0.5, 1.5] Hz and we observe an energy cascade up to 60 Hz. The theoretical slopes for gravity waves (≤ 15 Hz) and capillary waves predicted by the WTT are shown in red. We observe a spectrum much steeper than the theory in the capillary range. This is possibly due to dissipation, which becomes significant at these frequencies. Figure 3(b) shows the exponent extracted by fitting a power law to the temporal spectrum in the gravity range. For a large range of experiments we find an exponent near -5 with significant scattering. Although it is in contradiction of the theoretical prediction, it is actually in agreement with previous observations [13–16]. The scattering may be caused by the pollution of the water surface, which generates strong dissipation (by resonance between gravity-capillary waves and Marangoni waves [12]), although the experiments have been done with special care, i.e., filtered




FIG. 4. Space-time Fourier spectrum of the velocity field of the waves $E^v(k, \omega)$ for an experiment where (a) $\sigma = 0.025$ and (b) $\sigma = 0.08$. In both cases, the solid black line is the theoretical deep water linear dispersion relation for pure water $\omega^2 = gk + \frac{\gamma}{\rho}k^3$ with $\gamma = 72$ mN/m. The crossover between gravity and capillary waves occurs at $k_c = \sqrt{\rho g / \gamma} = 120\pi$ corresponding to a wavelength of 1.7 cm and a frequency equal to 13 Hz. Also shown are cross sections of $E^v(\mathbf{k}, \omega)$ at given frequencies at (c) weak forcing [$\sigma = 0.025$, same as (a)] and (d) stronger forcing. The solid black line is the linear dispersion relation.

water, careful cleaning, and protection of the wave tank, to limit the pollution by particles in the air. However, it remains difficult to maintain control of the surface quality over long times.

The full space-time measurements obtained with the Fourier transform profilometry allow us to compute the complete spectrum $E^v(\mathbf{k}, \omega)$. The computation of such space-time spectra is demanding in terms of data [notably for three-dimensional (3D) systems] but is an adequate tool to discriminate among various sorts of waves or nonpropagating motion [17–21]. The spectrum is computed as

$$E^v(\mathbf{k}, \omega) = \langle |v(\mathbf{k}, \omega)|^2 \rangle. \quad (5)$$

As for $E^\eta(\omega)$, the signal is divided into 16-s windows to compute the Fourier transform in time. In order to visualize $E^v(\mathbf{k}, \omega)$ on a 2D picture, we perform an angular integration of $\mathbf{k}$ to obtain $E^v(k, \omega)$. Figure 4(a) shows $E^v(k, \omega)$ at weak forcing. The energy is concentrated around the linear dispersion relation of gravity-capillary waves (3) (black line). The energy spread observed around the

dispersion relation is caused by weak nonlinear effects: The coherence of the linear waves is affected by the slow nonlinear energy exchanges [17,22] (note that an intrinsic finite spectral width exists also due to the finite spatial size of the images and to dissipation). The change of the width of the dispersion relation is thus a measure of the degree of nonlinearity of the system. Figure 4(c) shows four sections of the complete spectrum $E^v(\mathbf{k}, \omega)$ at given frequencies. We observe energy in almost all directions with a slight shift from the linear dispersion relation at high frequencies. This weak forcing corresponds to the second regime of turbulence reported by Cobelli *et al.* [13]. At increased forcing intensity [Figs. 4(b) and 4(d)], the spread of the energy around the dispersion relation is much wider, suggesting much stronger nonlinearities. Furthermore, a nonlinear shift is observable for all the frequencies. This nonlinear shift appears to be much stronger in the direction of the forcing Ox than in the transverse direction Oy . This is visible in the cross sections of the spectrum in Fig. 4(d) that are no longer circular as in Fig. 4(c). This deformation of the spectrum seems to be related to the amplitude of the forcing and is also qualitatively consistent with observations by Berhanu and Falcon [23], who interpret this shift as being due to nonlinear corrections due to large-amplitude low-frequency modes. They suggest that the dispersion relation could be

$$\omega^2 = \left\{ gk[1 + (ak)^2] + \frac{\gamma}{\rho}k^3 \left[1 + \left(\frac{ak}{4} \right)^2 \right]^{-1/4} \right\} \tanh(kh), \quad (6)$$

where a is the rms wave amplitude. If this effect actually depends on the propagation direction of the low-frequency Fourier modes, it could be responsible for the shift in the direction of the forcing for which the low-frequency components are stronger.

Thus our observations confirm the existence of two distinct regimes (weak and strong). The following section is devoted to a detailed study of the three-wave interaction that is supposed to be at the core of the energy cascade.

IV. THREE-WAVE RESONANCES

The most natural way to investigate nonlinear coupling among N waves is the use of N th-order correlations such as

$$\langle a_1 a_2 \cdots a_p a_{p+1}^* \cdots a_N^* \rangle, \quad (7)$$

where a_i is the wave amplitude in frequency space $a_i(\omega_i)$ or in wave-vector space $a_i(\mathbf{k}_i)$ and the asterisk denotes complex conjugation. This sort of generic correlation probes resonances $\omega_1 + \omega_2 + \cdots + \omega_p = \omega_{p+1} + \cdots + \omega_N$ or $\mathbf{k}_1 + \mathbf{k}_2 + \cdots + \mathbf{k}_p = \mathbf{k}_{p+1} + \cdots + \mathbf{k}_N$, respectively. In the following we investigate the three-wave case $N = 3$ in both frequency and wave-number space in order to get detailed information on the geometry of the wave interactions.

A. Theoretical analysis

The order of the nonlinear wave interaction depends on the order of the nonlinearity and the possibility to have solutions for the resonance equations. McGoldrick and then Simmons [8,9] investigated the three-wave resonant solutions in the gravity-capillary regime satisfying these two equations

$$\mathbf{k}_1 = \mathbf{k}_2 + \mathbf{k}_3, \quad (8)$$

$$\omega_1 = \omega_2 + \omega_3. \quad (9)$$

Figure 5(a) displays a full space-time representation of the resonant solutions for a given wave $[\mathbf{k}_2, \omega_2]$. The red surface shows the linear dispersion relation (3) and marks the point $[\mathbf{k}_1, \omega_1]$. As $\omega_1 = \omega_2 + \omega_3$, the green surface displays the dispersion relation shifted to start from the point $[\mathbf{k}_2, \omega_2]$ and thus marks the points $[\mathbf{k}_2 + \mathbf{k}_3, \omega_2 + \omega_3]$. Thus the resonant manifold is the intersection of the two surfaces marked by the thick black line. Due to the change of curvature of the dispersion relation, the intersection exists whatever the chosen value for ω_2 .




FIG. 5. Three-wave resonant solutions for gravity capillary waves. (a) Representation in 3D space. The red surface is the linear dispersion relation and a first wave $[\mathbf{k}_2, \omega_2]$ is given. A second dispersion relation is then plotted from this point (green surface). Thus, the resonant solution, satisfying Eqs. (8) and (9), is the intersection of these two surfaces (thick black line). A projection of the solutions in the $[k_x, k_y]$ space is plotted in red, showing then the possible wave-number $[\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3]$ combinations. (b) Plot of $\omega_3^{\min}$ as a function of ω_2 . Here $\omega_3^{\min}$ is the minimum value of ω_3 for which a resonant interaction is possible for a given ω_2 . (c) Corresponding value $\omega_1^{\min} = \omega_3^{\min} + \omega_2$.




FIG. 6. (a) Approximate resonant solution in the 1D case for a chosen k_2 (see the text). The dotted black lines represent a nonlinear widening of the dispersion relation by $\delta k/2\pi = 4 \text{ m}^{-1}$ [whose order of magnitude is estimated from $E^v(\mathbf{k}, \omega)$]. The green lines are the approximate solutions. (b) Color map of the distance Δk between the two dispersion relation [the black and red lines plotted in (a)] in the frequency space. The white line is the exact solution corresponding to $\Delta k/2\pi = 0 \text{ m}^{-1}$. The red line represents $\delta k/2\pi = 4 \text{ m}^{-1}$, which bounds the domain where approximate resonance occurs.

The resonant solutions show a point of minimum distance to the origin denoted by $[\mathbf{k}_3^{\min}, \omega_3^{\min}]$ that corresponds to the special 1D configuration where the three wave vectors $\mathbf{k}_1$, $\mathbf{k}_2$, and $\mathbf{k}_3$ are collinear. Figure 5(b) displays the variation of $\omega_3^{\min}$ as a function of ω_2 . In the whole area above this line exact resonant interactions are permitted. Thus, we observe the possibility of strong nonlocal coupling: A low-frequency gravity wave can interact only with capillary waves of much higher frequency. As it is well known, we see that pure gravity waves ($\omega/2\pi < 10 \text{ Hz}$) cannot interact with each other through a three-wave interaction. Figure 5(c) shows $\omega_1^{\min} = \omega_3^{\min} + \omega_2$ as a function of ω_2 to enhance the presence of the overall minimum of ω_1 . This special point corresponds to the degenerate case of Wilton waves: $\omega_1 = 2\omega_2 = 2\omega_3 = 2\pi \times 19.6 \text{ Hz}$ [Fig. 2(c)] [24]. The presence of such three-wave interactions has been verified experimentally by Henderson and Hammack [25].

As shown above in Fig. 4, nonlinear effects generate an energy spread around the dispersion relation. This may be seen as some uncertainty on the dispersion relation. We want to investigate the effect of this uncertainty on the resonances. For simplicity we consider only a unidirectional configuration of the wave vectors and we attribute all the (constant) uncertainty δk to the wave-vector equation. In this simplified case, Eq. (8) can be rewritten as

$$k_1 \pm \delta k = k_2 + k_3, \quad (10)$$

$$\omega_1 = \omega_2 + \omega_3. \quad (11)$$

Figure 6(a) shows the solution for a given wave $k_2/2\pi = 40 \text{ m}^{-1}$. The exact solution, similar to that in Fig. 5, corresponds to the intersection of the black and red curves. The nonlinear widening $\delta k/2\pi = 4 \text{ m}^{-1}$ [order of magnitude estimated from $E^v(\mathbf{k}, \omega)$ of the weak forcing in Fig. 4] is represented by the two dotted black lines near the thick black line of the dispersion relation. The range of new solutions permitted by this nonlinear widening are represented by the green lines. Figure 6(b) shows the distance Δk between the black and red lines as a function of frequency pairs (ω_2, ω_3) . The white line in Fig. 6(b) is the case of exact resonances for which the distance is zero. The thin red line limits the area in which the distance Δk is smaller than δk in which approximate




FIG. 7. Representation of three-wave coherence $C_\omega(\omega_1, \omega_2, \omega_3)$ where $\omega_2/2\pi = 10$ Hz (see the text for definitions) for (a) a weak forcing $\sigma = 0.025$ and (b) a strong forcing $\sigma = 0.08$. The surrounding blue background corresponds to statistical convergence noise so that the statistical convergence is about $10^{-2.5}$. The observed line of high correlation is above the convergence level by more than one order of magnitude at the maximum showing the presence of significant three-wave coupling in the signal. The line remains on the resonance curve $\omega_1 = \omega_2 + \omega_3$ as expected from such a statistical estimator for a stationary random signal. Correlations are $\log_{10}$ color coded.

resonances are possible. The range of possible resonances is thus strongly increased and thus these approximate resonances can increase significantly the intensity of energy transfer. In particular in a bounded system, where the discretization may not allow exact solutions, interactions may exist nonetheless. Note that approximate resonances also allow three-wave interactions in the gravity regime, whereas exact resonances are only possible among four waves. Indeed, as the two dispersion relations are very close for these gravity waves, a weak nonlinear widening permits three-wave interactions. This analysis could be extended to the 2D case (Fig. 5) and we could also make δk vary as a function of ω to be closer to the experiments, but in both cases we do not expect a significant change from the simple 1D case.

B. Frequency correlations

As we expect three-wave resonances, we focus our study on three-wave correlations. The simplest case is to work in the frequency space due to the reduced number of parameters: three frequencies reduced to two using the resonant equation (9). From $v(x, y, t)$ we compute the Fourier transform in time over 4-s time windows so as to obtain $v(x, y, \omega)$. Correlations are then computed as

$$C_\omega(\omega_1, \omega_2, \omega_3) = \frac{|\langle\langle v^*(x, y, \omega_1)v(x, y, \omega_2)v(x, y, \omega_3) \rangle\rangle|}{[E^v(\omega_1)E^v(\omega_2)E^v(\omega_3)]^{1/2}}, \quad (12)$$

where $\langle\langle \dots \rangle\rangle$ denotes a double average: a space average over (x, y) on the image and over successive time windows. Here $E^v(\omega) = \langle\langle |v(x, y, \omega)|^2 \rangle\rangle$ is the frequency spectrum. The coherence defined in this way should be zero for uncorrelated waves and one for perfectly correlated waves. For a stationary random signal such a correlation is expected to be nonzero only along the resonance line $\omega_1 = \omega_2 + \omega_3$. Figure 7 shows the coherence map at a given frequency $\omega_2/2\pi = 10$ Hz. We observe for both forcing intensities that a line of correlation emerges from the statistical convergence noise




FIG. 8. Bicoherence $B_\omega(\omega_2, \omega_3)$ for (a) a weak forcing where $\sigma = 0.02$ and (b) a strong forcing where $\sigma = 0.08$. The solid black line represents the border $\omega_3^{\min}$ of the area of possible exact resonant solutions. The dotted black lines represent the 1D solution affected by a nonlinear spectral spreading $\delta k/2\pi = 4 \text{ m}^{-1}$. Correlations are $\log_{10}$ coded.

that is less than 3×10^{-3} . This implies the presence of a significant three-wave resonant process in the signal. We note a difference between the two forcing intensities: At weak forcing a correlation peak is seen near 20 Hz, whereas the strong forcing appears much more homogeneous in terms of variation of the correlation amplitude with the frequency. For exact resonances, with a given value of $\omega_2 = 10 \text{ Hz}$, the minimum value of ω_3 that is allowed is also 10 Hz and the corresponding minimum value of ω_1 is about 20 Hz (see Fig. 5). Thus, at the weakest forcing one expects that the correlation becomes nonzero only for values of ω_1 larger than 20 Hz. The fact that the correlation is weak for values of ω_1 larger than 20 Hz suggests that the unidirectional coupling (corresponding to $\omega_1 = 20 \text{ Hz}$) is dominant. At larger forcing intensity the approximate resonances become numerous and thus the wave coupling involves many more wave triads.

In order to perform a deeper analysis of the resonant interaction, we compute the bicoherence, defined as

$$B_\omega(\omega_2, \omega_3) = C(\omega_2 + \omega_3, \omega_2, \omega_3). \quad (13)$$

This corresponds to the extraction of the resonant line observed in Fig. 7 for all frequencies ω_2 . The presence of this converged line above the noise level has been checked for all frequencies. Figure 8 shows the bicoherence map at the two forcing amplitudes. In the weak forcing case [Fig. 8(a)], a noticeable organization of the coherence can be seen. First, the figure is symmetric along the diagonal $\omega_2 = \omega_3$ as expected from the symmetry of the resonance relations when permuting the indices $2 \leftrightarrow 3$. Second, a wide line of high coherence (about 10^{-1}) is located close to the $\omega_3^{\min}$ curve (thick black line) described previously and corresponding to exact resonances in the unidirectional case. In the region of allowed exact resonances (above the thick black line) the coherence level above this curve remains low ($\lesssim 10^{-2}$). This suggests that the dominant interactions are close to unidirectional. In this case the observed isotropy of the spectrum could be due mostly to the fact that the curved walls create a chaotic cavity. Nevertheless, experiments performed in a rectangle cavity do not show much difference in the isotropy. This is most likely due to the fact that the coupling still permits some angular spreading or energy, especially at low wave number, as confirmed by the wave-number resonance analysis below.

At very low values of both ω_2 and ω_3 a few spots of high coherence exist that correspond to the allowed approximated three-wave resonances among gravity waves. We observe another region of high coherence for small values of ω_3 and large values of ω_2 [bottom part of Fig. 8(a)]. This region is composed of two strong lines of high coherence, corresponding to the frequencies of the first eigenmodes of the wave tank. We can explain the existence of this region with the approximate resonances discussed above. As the interactions are observed to be mostly one dimensional, we show as dotted black lines the region of approximate resonances in the 1D case for $\delta k/2\pi = 4 \text{ m}^{-1}$, already shown in Fig. 6(b). This solution encircles very convincingly the high correlation area and highlights the importance of these approximate resonances: The number of effective interactions increases significantly and they do operate. The case of stronger forcing is displayed in Fig. 8(b). The coherence is less organized and less straightforward to interpret. The maximum level of correlation is of the same order as the weak forcing, but the 1D resonant line is less visible. The spread of the high coherence region is much wider than in the low forcing case. The coherence level is much higher also in the regions that appear in blue in Fig. 8(a). Wave interactions seem possible in all the domain and there are no longer any indications of a possibly dominant unidirectional process. We also note that some discretization become apparent at higher frequencies. It is consistent with the peaks on the temporal spectrum visible in Fig. 3(a). These peaks are most likely related to the strong forcing, although their interpretation is not fully clear so far.

C. Space analysis

In order to confirm directly the unidirectional character of the interactions, we investigate the correlations in the wave-number domain. Figure 9 shows the expected exact solutions projected in $[k_x, k_y]$ space. It is a more detailed view of the projections of the solution shown in Fig. 5(a). Figure 9(a) illustrates the method to find the resonances: For simplicity we choose the 1D case at a given $\mathbf{k}_2$ with the inclusion of the negative frequencies and wave numbers. The black lines are the dispersion relations and the red lines are the shifted dispersion relations as in Sec. IV A. The resonant solutions are thus the intersections of the black and red curves and they are marked as green dots.

Figure 9(b) represents the general 2D solutions in $[k_x, k_y]$ space for a given wave vector $\mathbf{k}_2$. In this representation, exact 1D solutions appear as points and are reported as the green dots. The arrows show an example of the allowed wave-number combination for the given vector $\mathbf{k}_2$ (in red). The symmetry in frequency implies that solutions A1 and A2 (as for A3 and A4) are superposed in $[k_x, k_y]$ space. To go further, the two branches of solutions are equivalent and correspond to permutations of indices in the resonant equations. For instance, solution A1 corresponds to $\omega_1 = \omega_2 + \omega_3$ and solution A4 corresponds to $\omega_1 = -\omega_2 + \omega_3$, which can be rewritten as $\omega_3 = \omega_1 + \omega_2$. Thus, by permuting the indices, they are fundamentally equivalent. Figures 9(c) and 9(d) show the apparition of a new solution at larger k_2 . As we have just seen previously, they are also equivalent by permutation of the indices.

Similarly to the definition of $C_\omega(\omega_1, \omega_2, \omega_3)$ in the frequency case, we define the correlations in space $C_k(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3)$ as

$$C_k(\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}_3) = \frac{|\langle v^*(\mathbf{k}_1, t)v(\mathbf{k}_2, t)v(\mathbf{k}_3, t) \rangle|}{[E^v(\mathbf{k}_1)E^v(\mathbf{k}_2)E^v(\mathbf{k}_3)]^{1/2}}. \quad (14)$$

Here the average stands for an average over time and $E^v(\mathbf{k}) = \langle |v(\mathbf{k}, t)|^2 \rangle$ is the spatial Fourier spectrum. We thus define the bicoherence $B_k(\mathbf{k}_2, \mathbf{k}_3)$ as

$$B_k(\mathbf{k}_2, \mathbf{k}_3) = C(\mathbf{k}_2 + \mathbf{k}_3, \mathbf{k}_2, \mathbf{k}_3). \quad (15)$$

Unlike the frequency correlation, $B_k(\mathbf{k}_2, \mathbf{k}_3)$ is a function of four real parameters, which makes its representation more difficult. In order to obtain a 2D picture that is easier to analyze, we first choose a value of $|\mathbf{k}_2| = k_2$. We assume that the turbulent wave field is statistically isotropic and we perform an average over eight distinct directions, after rotation of the whole image so that $\mathbf{k}_2$ lies on the




FIG. 9. (a) Theoretical exact solutions in the unidirectional case for a given wave number $k_2/2\pi = 50 \text{ m}^{-1}$. The black lines are the dispersion relations and the red lines are the shifted dispersion relations as in Sec. IV A; the resonant waves are found at their intersection (green dots). (b) Theoretical exact solutions in the general case, in (k_x, k_y) space and for a given wave vector of modulus equal to $k_2/2\pi = 50 \text{ m}^{-1}$. Exact 1D solution corresponding to (a) are reported by the green points. Arrows represent a resonant triplet. (c) One-dimensional theoretical exact solutions for a larger wave number $k_2/2\pi = 110 \text{ m}^{-1}$. We observe the apparition of a new region of solutions in addition to the previous observed in (a). This new solution is actually equivalent to the other by permutation of the indices of the waves. (d) Two-dimensional theoretical exact solutions in (k_x, k_y) space for a given $k_2/2\pi = 110 \text{ m}^{-1}$.

horizontal axis. This permits us to compute $B_k(k_2, \mathbf{k}_3)$ with a better statistical convergence. Figure 10 displays $B_k(k_2, \mathbf{k}_3)$ for three k_2 and two forcings (weak on the left, strong on the right). The exacts solutions are plotted in black and the given $\mathbf{k}_2$ are represented by the black arrows. We observe a concentration of high correlation along the exact solutions presented before. As expected, due to the permutation equivalence, there is a vertical symmetry axis at $k_x/2\pi = 0$ and $k_x/2\pi = k_2/2$. The




FIG. 10. Bicoherence $B_k(k_2, \mathbf{k}_3)$ for $\sigma = 0.02$ (left) and $\sigma = 0.08$ (right). We fixed (a) and (d) $|\mathbf{k}_2|/2\pi = 20$, (b) and (e) $|\mathbf{k}_2|/2\pi = 50$, and (c) and (f) $|\mathbf{k}_2|/2\pi = 125$. Correlations are averaged over eight different directions for $\mathbf{k}_2$ (after rotation of the whole picture). The black arrow represents the chosen $\mathbf{k}_2$ and the exact solution is plotted in black. Color is $\log_{10}$ coded. We observe a concentration of correlation emerging from the statistical noise along the exact solutions. The spreading along the line comes from the approximate resonances. The presence of a maximum along the direction of $\mathbf{k}_2$ confirms the dominance of the unidirectional interactions. For low $\mathbf{k}_2$ we note a significant angular spreading of interactions.

magnitude of the correlation is about the same level of 10^{-1} observed for the frequency bicoherence $B_\omega(\omega_2, \omega_3)$. For the weak forcing, the repartition of the correlations shows a concentration of higher value along the 1D direction (horizontal axis with $k_y = 0$), consistently with our conclusions from




FIG. 11. Linear phase velocity $v_\phi = \frac{\omega}{k}$ and the linear group velocity $v_g = \frac{\partial \omega}{\partial k}$ in the conditions of our experiment. Here v_ϕ gives a minimum near 15 Hz, where the waves are nondispersive. We note also that v_ϕ is quasiconstant on the interval $[10, 30]$ Hz.

the analysis of $B_\omega(\omega_2, \omega_3)$. However, the observed resonances allow for a significant coupling of wave vectors with a quite large angle among them, notably at small k_2 . The stronger forcing presents the same general behavior with a much wider nonlinear spreading resulting, as expected, by the approximate resonances. The observed correlations also allow a much higher coupling between wave vectors that are not collinear.

V. CONCLUSION

The investigation of three-wave correlations confirms the existence of three-wave nonlinear coupling in weak wave turbulence. In the weakest case, the observed correlations suggest that the dominant coupling occurs for wave triads that are close to collinear. If the dispersion relation takes the form of a pure power law $\omega \propto k^\alpha$ (with $\alpha \neq 1$ for dispersive waves), no three-wave resonance is possible for collinear wave vectors. In the case of gravity-capillary waves, the change of curvature allows for such resonances if waves have sufficiently separated frequencies. Our observations also underline the importance of approximate resonances permitted by nonlinear spectral widening. These approximate resonances permit notably three-wave interactions among gravity waves that may induce a more efficient energy transfer among gravity waves than the four-wave exact resonances.

The fact that the unidirectional resonances seem to dominate the correlations may be related to the fact that the phase and group velocities exhibit a minimum around the frequency of the gravity-capillary crossover. The linear phase velocity $v_\phi = \frac{\omega}{k}$ and the linear group velocity $v_g = \frac{\partial \omega}{\partial k}$ are shown in Fig. 11. A direct effect of this double curvature is the presence of a local minimum on the phase velocity near $\omega/2\pi = 15$ Hz and a minimum of the group velocity around 8 Hz. In addition, we observe that v_ϕ remains roughly constant near this minimum in the interval $[10, 30]$ Hz and the group velocity increases very slowly in the same range of frequencies: These waves are thus weakly dispersive. The combination of these two effects may increase significantly the efficiency of the 1D interaction. Indeed, the waves packets moving with similar speeds in the same direction have a long time to interact. Then, for a weakly nonlinear system, where the interaction time has to be long, 1D interactions are enhanced and finally dominate. When the degree of nonlinearity increases, the interaction time becomes shorter, thus unidirectional interactions may not dominate

as significantly. Thus, waves traveling with a slight angle may have time to interact and exchange similar amounts of energy. We also observed that the maximum level of correlation is constant for all our experiments (about 10^{-1}). This constant value of the correlation suggests that the efficiency of the nonlinear interactions may be bounded. However, this possible saturation of the interactions remain to be confirmed with more strongly nonlinear experiments.

The weak turbulence theory has been applied separately both to pure gravity surface waves and to capillary waves. Our observations show that at the crossover between both types of waves, the wave turbulence that is observed may be quite far from what is expected from the study of either regime separately. The striking example is the possibility to observe weak nonlinear interactions among collinear waves that may even be dominant due to the low dispersion of the waves in this range of frequencies. Another feature is the possibility to observe nonlocal coupling between gravity waves and capillary waves as well as three-wave coupling of gravity waves allowed by nonlinear spectral widening. Our observations illustrate the importance of approximate resonances permitted by nonlinearity that may change very significantly the dynamics of wave turbulence as compared to the case of exact resonances. If three-wave coupling is permitted we should expect it to be much more efficient to transfer energy than the four-wave coupling predicted in the case of exact resonances. It remains to investigate whether this three-wave coupling is indeed observed in oceanic data. Another question is also the range of frequencies or wavelengths of gravity waves that may be impacted by this mechanism. Is it restricted to frequencies close to the gravity-capillary crossover or does it have an impact on much lower frequencies, for long waves?

ACKNOWLEDGMENTS

This project received funding from the European Research Council under the European Union's Horizon 2020 research and innovation programme (Grant No. 647018-WATU). We thank Kronos Worldwide, Inc. for kindly providing us with the titanium oxide pigment Kronos 1001.

-
- [1] V. E. Zakharov, V. S. L'vov, and G. Falkovich, *Kolmogorov Spectra of Turbulence* (Springer, Berlin, 1992).
 - [2] S. Nazarenko, *Wave Turbulence* (Springer, Berlin, 2011).
 - [3] A. C. Newell and B. Rumpf, Wave turbulence, [Annu. Rev. Fluid Mech.](#) **43**, 59 (2011).
 - [4] E. A. Kartashova, Weakly Nonlinear Theory of Finite-Size Effects in Resonators, [Phys. Rev. Lett.](#) **72**, 2013 (1994).
 - [5] E. Kartashova, Model of laminated wave turbulence, [JETP Lett.](#) **83**, 283 (2006).
 - [6] E. Kartashova, Discrete wave turbulence, [Europhys. Lett.](#) **87**, 44001 (2009).
 - [7] Y. V. Lvov, S. Nazarenko, and B. Pokorni, Discreteness and its effect on water-wave turbulence, [Physica D](#) **218**, 24 (2006).
 - [8] L. F. Mcgoldrick, Resonant interactions among capillary-gravity waves, [J. Fluid Mech.](#) **21**, 305 (1965).
 - [9] W. F. Simmons, A variational method for weak resonant wave interactions, [Proc. R. Soc. London Ser. A](#) **309**, 551 (1969).
 - [10] Q. Aubourg and N. Mordant, Nonlocal Resonances in Weak Turbulence of Gravity-Capillary Waves, [Phys. Rev. Lett.](#) **114**, 144501 (2015).
 - [11] P. J. Cobelli, A. Maurel, V. Pagneux, and P. Petitjeans, Global measurement of water waves by Fourier transform profilometry, [Exp. Fluids](#) **46**, 1037 (2009).
 - [12] A. Pradka, B. Cabane, V. Pagneux, A. Maurel, and P. Petitjeans, Fourier transform profilometry for water waves: How to achieve clean water attenuation with diffusive reflection at the water surface? [Exp. Fluids](#) **52**, 519 (2012).
 - [13] P. Cobelli, A. Pradka, P. Petitjeans, G. Lagubeau, V. Pagneux, and A. Maurel, Different Regimes for Water Wave Turbulence, [Phys. Rev. Lett.](#) **107**, 214503 (2011).
 - [14] S. Nazarenko, S. Lukaschuk, S. McLelland, and P. Denissenko, Statistics of surface gravity wave turbulence in the space and time domains, [J. Fluid Mech.](#) **642**, 395 (2009).

- [15] B. Issenmann and E. Falcon, Gravity wave turbulence revealed by horizontal vibrations of the container, [Phys. Rev. E **87**, 011001 \(2013\)](#).
- [16] L. Deike, B. Miquel, P. Gutierrez, T. Jamin, B. Semin, M. Berhanu, E. Falcon, and F. Bonnefoy, Role of the basin boundary conditions in gravity wave turbulence, [J. Fluid Mech. **781**, 196 \(2015\)](#).
- [17] N. Mordant, Fourier analysis of wave turbulence in a thin elastic plate, [Eur. Phys. J. B **76**, 537 \(2010\)](#).
- [18] E. Herbert, N. Mordant, and E. Falcon, Observation of Nonlinear Dispersion Relation and Spatial Statistics of Wave Turbulence on the Surface of a Fluid, [Phys. Rev. Lett. **105**, 144502 \(2010\)](#).
- [19] P. Clark di Leoni, P. Cobelli, and P. D. Mininni, The spatio-temporal spectrum of turbulent flows, [Eur. Phys. J. E **38**, 1 \(2015\)](#).
- [20] P. Clark di Leoni, P. D. Mininni, and M. E. Brachet, Spatiotemporal detection of Kelvin waves in quantum turbulence simulations, [Phys. Rev. A **92**, 063632 \(2015\)](#).
- [21] F. Leckler, F. Ardhuin, C. Peureux, A. Benetazzo, F. Bergamasco, and V. Dulov, Analysis and interpretation of frequency—Wavenumber spectra of young wind waves, [J. Phys. Oceanogr. **45**, 2484 \(2015\)](#).
- [22] B. Miquel and N. Mordant, Nonlinear dynamics of flexural wave turbulence, [Phys. Rev. E **84**, 066607 \(2011\)](#).
- [23] M. Berhanu and E. Falcon, Space-time-resolved capillary wave turbulence, [Phys. Rev. E **87**, 033003 \(2013\)](#).
- [24] J. Wilton, LXXII. On ripples, [London Edinburgh Dublin Philos. Mag. J. Sci. **29**, 688 \(1915\)](#).
- [25] D. Henderson and J. Hammack, Experiments on ripple instabilities. Part 1. Resonant triads, [J. Fluid Mech. **184**, 15 \(1987\)](#).